

Neotropical Diptera

Neotropical Diptera 11: 1-17 (April 15, 2009)
ISSN 1982-7121
www.neotropicaldiptera.org

Depto. de Biologia – FFCLRP
Universidade de São Paulo
Ribeirão Preto, SP, Brazil

Catalogue of Neotropical Diptera. Cuterebridae¹

Nelson Papavero

Museu de Zoologia, Universidade de São Paulo, São Paulo, SP, Brasil
Pesquisador Visitante do Departamento de Biologia, Faculdade de Filosofia, Ciências e Letras, Universidade de São Paulo, Ribeirão Preto, SP, Brasil

&

†José Henrique Guimarães

Departamento de Parasitologia, Instituto de Ciências Biomédicas
Universidade de São Paulo
São Paulo, SP

Introduction

In the region covered by this catalogue, the Cuterebridae are represented by six genera and 37 species, as shown in the table below, plus eight unrecognized or unidentified species, and two *nomina nuda*.

Genus	Number of species
<i>Andinocuterebra</i>	1
<i>Cuterebra</i>	16
<i>Dermatobia</i>	1
<i>Metacuterebra</i>	14
<i>Pseudogametes</i>	2
<i>Rogenhofera</i>	5

Family Cuterebridae Brauer, 1863

Cuterebriden (vernacular) Brauer, 1863b: 64, 70. Type-genus, *Cuterebra* Clark, 1815.

Cuterebridae Brauer, 1887: 5.

Cutiterebrinae Austen, 1895a: 378.

Cuterebrinae Bau, 1906: 24 (as subfamily of "Muscaridae").

Pseudogametinae [sic] Townsend, 1911a: 139. Type-genus, *Pseudogametes* Bischoff, 1900.

Pseudogametini Townsend, 1911b: 329 (correction of ending without naming).

Cuterebrinae Townsend, 1913: 57.

¹ This project was supported by FAPESP grants # 2003/10.274-9, 2007/50877-5, and 2007/50878-1.

Dermatobiiae Townsend, 1913: 57. Type-genus, *Dermatobia* Brauer, 1861.
Dermatobiinae Townsend, 1917: 23.
Pseudogametinae Townsend, 1917: 23.
Cuteribrinae Rodhain, 1927: 200.
Dermatobidae Bromley, 1937: 105.
Pseudomagetini Dalmat, 1942a: 23, 24, error.
Rogenhoferinae J. H. Guimarães, 1990: 255. Type-genus, *Rogenhofera* Brauer, 1863.

Genus **Andinocuterebra** J. H. Guimarães, 1984

Andinocuterebra J. H. Guimarães, 1984: 23. Type-species, *Andinocuterebra fassleri* J. H. Guimarães, 1984 (orig. des.).

fassleri J. H. Guimarães, 1984: 23. Type-locality: Colombia, Huila, río Aguacatal. Distr. – Colombia (Huila). Refs. – J. H. Guimarães, 1990: 261; J. H. Guimarães & Papavero, 1999: 202, fig. 96.

Genus **Cuterebra** Clark, 1815

Cuterebra Clark, 1815: 64, 70. Type-species, *Oestrus cuniculi* Clark, 1797 (sub. des., Desmarest, 1859: 247, as “*Cuterebra cuniculi* Fabr.”).

Trypoderma Wiedemann in Meigen, 1824: 165. Type-species, *Musca americana* Fabricius, 1775 (sub. des., Wiedemann, 1830: 256).

Curtebra Blanchard, 1840: 607, error.

Curtterebra Blanchard, 1845: 484, error.

Cuttiterebra Agassiz, 1846: 108, unjust. emend.

Cutirebra Wulp, 1888: 1, error.

Bogeria Austen, 1895a: 391. Type-species, *Bogeria princeps* Austen, 1895 (orig. des.).

Atrypoderma Townsend, 1919: 592. Type-species, *Musca americana* Fabricius, 1775 (orig. des.).

Cuterebra, subg. *Orthocuterebra* Bau, 1929b: 542. Type-species, *Cuterebra lepusculi* Townsend, 1897 (sub. des., Aldrich, 1931: 117).

Cuterebra, subg. *Paracuterebra* Bau, 1929b: 543. Type-species, *Oestrus cuniculi* Clark, 1797 (sub. des., Aldrich, 1931: 117).

Cuterebra, subg. *Protocuterebra* Bau, 1929b: 543. Type-species, *Musca americana* Fabricius, 1775 (sub. des., Aldrich, 1931: 117).

Atryposoma Bau, 1931a: 20, error.

Pseudobogeria Bau, 1931b: 206. Type-species, *Oestrus buccatus* Fabricius, 1776 (orig. des.).

Cuterebrella J. H. Guimarães, 1984: 24, figs. 2-3. Type-species, *Cuterebra praegrandis* Austen, 1933 (orig. des.). Refs. – Sabrosky, 1986: 22 (as ?syn.); J. H. Guimarães & Papavero, 1999: 203 (syn.).

Refs. – Dixon, Winkler & Nelson, 1969 (ophthalmomyiasis); Rice & Douglas, 1972 (human myiasis); Rodrigues, Weiss & Muncy, 1974 (ophthalmomyiasis); Scholten & Chrom, 1979 (human myiasis); Timm & Lee Jr., 1981 (effects on hosts); Baird, Podgore & Sabrosky, 1982 (myiasis in humans); Sabrosky, 1986: 43-55 (key to North and part of Central American spp.); Baird, Baird & Sabrosky, 1989 (human myiasis); Schiff, 1993 (furuncular cutaneous myiasis); Glasgow & Maggiano, 1995 (ophthalmomyiasis); Glass, Cornetta, Lahunta, Center & Kent, 1998 (myiasis of central nervous system in cats); J. H. Guimarães & Papavero, 1999: 203; Keth, 1999 (human myiasis); Cornet, Florent, Lefebvre, Wertheimer, Perez-Eid, Bangs & Bouvet, 2003 (tracheopulmonary myiasis, review).

albipilosa Sabrosky, 1986: 177. Type-locality: U. S. A., California, San Diego County, Summit Cowles Mt. Distr. – U. S. A. (Arizona, California, Colorado, New Mexico), Mexico (Baja California Sur).

arizonae Sabrosky, 1986: 186, map. 16. Type-locality: U. S. A., Arizona, Santa Cruz County, Madera Canyon, Santa Rita Mts. Distr. – U. S. A. (Arizona), Mexico (Sonora).

atrox Clark, 1848: unnumb. page, fig. 5. Type-locality: “Africa” [in error, actually Mexico]. Distr. – Southwestern U. S. A.

(Arizona, Colorado, New Mexico, western Texas), and widespread in Mexico (Chihuahua to Nuevo León, south to Chiapas). Refs. – Brauer, 1863b: 229; Wulp, 1888: 2; Austen, 1895a: 378 (*Cutiterebra*), 1895b: 147 (*Cutiterebra*); Aldrich, 1905: 418 (cat.); Swenk, 1905: 182 (in key); Jones, 1906: 392 (in key); Townsend, 1927: 24; Bau, 1928a: 51 (in key), 1929a: 3 (in key), 1929b: 544 (*Cuterebra (Oricuterebra)*), 1931: 207 (*Cutrerebra (Atrypoderma)*); Austen, 1933: 701; Townsend, 1935: 220 (*Atrypoderma*); J.H. Guimarães, 1967: 2 (cat.); Sabrosky, 1986: 148, map 11, figs. 39-41; Papavero & Ibáñez Bernal, 2001: 89 (hist. Mexican dipterol.).

similis Johnson, 1903: 101. Type-locality: U. S. A., New Mexico, Beulah.

americana Fabricius of Johnson, 1903: 105, misident.

austeni Sabrosky, 1986: 189, map 17. Type-locality: U. S. A., Arizona, Pinal County. Distr. – U. S. A. (Arizona), Mexico (Sonora). Refs. – Baird, 1997: 690 (biol.); Wilson, Hnida & Duszynski, 1997: 359 (hosts); Colwell, Baird, Lee & Milton, 1999: 803 (SEM and morphometrics of egg); Kemp & Alcock, 2003: 290 (biol.); Alcock & Kemp, 2004: 273 (mating system).

bajensis Sabrosky, 1986: 163, map 13. Type-locality: Mexico, Baja California Norte, 8 km north of Punta Prieta. Distr. – U. S. A. (California), Mexico (Baja California Norte, Baja California Sur).

buccata (Fabricius), 1777: 305 (*Oestrus*). Type-locality: U. S. A., “Carolina” [i. e., South Carolina]. Distr. – U. S. A. (North Dakota to New Hampshire (or to Nova Scotia?), south to Texas and Florida, also New Mexico, Arizona), Mexico (Chiapas), Guatemala and Honduras. Refs. – Fabricius, 1781: 398 (*Oestrus*), 1787: 305 (*Oestrus*), 1794: 230 (*Oestrus*); Latreille, 1803: 50 (*Oestrus*); Fabricius, 1805: 227 (*Oestrus*); Olivier, 1812: 464 (*Oestrus*); Latreille, 1818: 270 (n. comb.); Meigen, 1824: 165 (*Trypoderma*); Wiedemann, 1830: 259 (*Trypoderma*); Macquart, 1835: 47; Harris, 1841: 420 (*Oestrus*); Brauer, 1863b: 230 (in key); Austen, 1895a: 384 (*Cutiterebra*); Johnson, 1895: 331; Johnson, in J. B. 1900: 667; Swenk, 1905: 182 (in key); Washburn, 1905: 138, fig. 130; Johnson in J. B. Smith, 1910: 774; Johnson, 1913: 69; Townsend, 1917: 24 (in key; *Bogeria*); Johnson, 1925a: 182 (*Bogeria*), 1925b: 91 (*Bogeria*); Bau, 1929a: 4 (in key), 1929b: 544 [*Cuterebra (Orthocuterebra)*], 1929c: 305, fig. 7; Painter, 1930: 32 (*Bogeria*); Bau, 1931b: 207 [*Cuterebra (Pseudobogeria)*]; Davis, 1937: 85; Knipling & Bruce, 1937: 48; Lindquist, 1937: 736; Brimley, 1938: 357; Townsend, 1938: 172 (*Pseudobogeria*); Knipling & Brody, 1940: 33, pls. 1-2; Dalmat, 1942c: 420, fig.; Haugen, 1942: 229; Dalmat, 1943a: 183; Bequaert, 1945: 176 (human myiasis); Jellison, 1949: 485; Beamer, 1950: 16; Penner, 1958: 67, figs. (of egg and 1st and 2nd instar larvae); Zimsen, 1964: 478 (*Oestrus*); Cole, 1970: fig. 355 (p. 585); Sabrosky, 1972: 94; Weisbroth, Wang & Scher, 1973: 22 (immune response in myiasis of domestic rabbits); Jacobson, McGinnes & Catts, 1978: 56 (biol.); Magnarelli & Andreadis, 1981: 894 (human myiasis); Sabrosky, 1986: 119, map 6, figs. 35-36; Nelder & Reeves, 2005: 316; Slansky, 2006: 154 (hosts).

purivora Clark, 1815: 70, pl. 2, fig. 29. Type-locality: “America”. Ref. – Sabrosky, 1972: 94 (LT selection].

cuccatus Harris, 1833: 594 (1835: 599), error.

?*cuniculi* Clark of Harris, 1833: 594 (1835: 599) and R. C. Smith, 1943: 374, misident.

horripilum Clark of Walker, 1849: 683, misident.

buccator Williston, 1884: 426, fig. 536, error.

baccata Lugger, 1896: 186, 1897: 229, fig. 186, error.

bacata Washburn, 1905: 137, error.

scudderi Townsend, 1917: 24 (in key), 27. Lectotype-locality: U. S. A., Washington, D. C. Refs. – Roberts, 1933: 159. sp. Curran, 1934: 411, fig.

buccatta Eddy & Emerson, 1940: 45, error.

clarki Sabrosky, 1986: 145, map 10. Type-locality: Mexico, Nuevo León, El Diente, just outside Monterrey. Distr. – U. S. A. (Oklahoma), Mexico (Nuevo León, Tamaulipas).

fontinella Clark, 1827: 410. Lectotype-locality: U. S. A., Illinois. Distr. – Southern Canada, U. S. A. (except Alaska and Hawaii), Mexico (Nuevo León, Tamaulipas). Refs. – Clark, 1815: pl. 2, fig. 23 (unnamed); Joly, 1846: 289; Brauer, 1863b: 230 (in key), 242; Austen, 1895a: 384, 1895b: 152 (*Cutiterebra*); Coquillett, 1898: 9; Swenk, 1905: 182; Johnson, 1913: 70; Townsend, 1917: 24 (*Bogeria*); Cole & Lovett, 1921: 298; Johnson, 1925a: 182 (*Bogeria*); Bequaert, 1928: 805 (*Bogeria*); Bau, 1929a: 5 (in key), 1929b: 544 [*Cuterebra (Orthocuterebra)*]; Johnson, 1930: 283; Bau, 1931: 207 [*Cuterebra (Pseudobogeria)*]; Austen, 1933: 705; Greene, 1935 (biol.); Townsend, 1938: 173 (*Psaeudobogeria*); Brimley, 1938: 357; Burt, 1940: 41 (as *Cuterebra* larvae; biol.); Scott & Smead, 1942: 94; Dalmat, 1942b: 47; Penner & Pocius, 1956 (biol.); Sillman, 1956, 1957a, 1957b (biol.); Miller & Getz, 1969 (biol.); Drummond, Gingrich & Gladney, 1970: 85 (used in tests for systemic insecticide); Sabrosky, 1972: 94 (LT des.); Gingrich, 1975: 67 (larval migration behavior in *Peromyscus leucopus*); Hensley, 1976: 172; Gingrich, 1979: 288; Parker & Chaney, 1979 (biol.); Timm & Cook, 1979: 211 (effect on reproduction in *Peromyscus leucopus*); Parker & Chaney, 1979: 573; Catts, 1982 (biol.); Pruitt & Barrett, 1983: 113; Shiffer, 1983 (aggregation behavior); Goddard, Baker & Norment, 1986: 128; Sabrosky, 1986: 70, map 2, frontispiece, figs. 13, 20-21; Cogley, 1991: 275; Durden, 1995: 787; Colwell, Baird, Lee & Milton, 1999: 803 (SEM and morphometrics of egg); Wolf & Batzli, 2001: 106; Noël, Tessier, Angers, Wood & Lapointe, 2004: 161; Cramer & Cameron, 2006: 1103; Jennison, Rodas & Barrett, 2006:

- 157; Slansky, 2006: 154 (hosts); Slansky & Huckabee, 2006: 1369 (in *Procyon lotor* (Carnivora, Procyonidae)); Cramer & Cameron, 2007: 1419; Barrett & Feldhamer, 2008.
- fontanella* Clark, 1843: 88 and Walker, 1849: 100, error.
- americana* Fabricius of Walker, 1849: 683, misident.
- cuniculi* Clark of Johnson, 1895: 331, misident.
- frontinella* Fletcher & Gibson, 1908: 38, error.
- sp.* Townsend, 1911a: 137 (descr. of egg).
- buccata* Fabricius of Johnson, 1913: 69, misident.
- fasciata* Swenk of Townsend, 1917: 24 (*Bogeria*), and Cameron, 1926: 430, misident.
- peromysci* Dalmat, 1942b: 45 (descr., 3 larval stages, puparium). Type-locality: U. S. A., Iowa, Ames. Ref. – Dalmat, 1942c: 419, 1943: 311; Jellison, 1949: 485.
- angustifrons* Dalmat, 1942c: 418. Type-locality: U. S. A., Iowa, Ames. Ref. – Sillman, 1957a: 28 (ecol., life history); Sillman & Smith, 1959: 165 (exp. infestation of *Peromyscus leucopus* with larvae); Shoemaker & Joy, 1968: 78; Getz, 1970: 187; Sabrosky, 1972: 94 (syn.); Hunter & Webster, 1973: 1301.
- sp.* Sillman, 1956: 89.
- histrio** Coquillett, 1902: 103. Type-locality: Mexico, Guanajuato. Distr. – Mexico (Chihuahua, Durango, Guanajuato, San Luis Potosí, Veracruz, Zacatecas). Refs. – Swenk, 1905: 182 (in key); Townsend, 1917: 24 (in key), 25; Bau, 1929a: 4 (in key), 1929a: 544 [*Cuterebra (Orthocuterebra)*], 1931a: 20 (fig. 2) [*Cuterebra (Cuterebra)*], 1931b: 205 [*Cuterebra (Cuterebra)*]; J.H. Guimarães, 1967: 3 (cat.); Sabrosky, 1986: 168, map 13, fig. 44.
- americana* Fabricius of Wulp, 1888: 1, part, misident.
- lepusculi** Townsend, 1897: 8. Type-locality: U. S. A., New Mexico, Doña Ana. Distr. – U. S. A. Washington to South Dakota, south to California and Texas), Mexico (Morelos). Refs. – Swenk, 1905: 182 (in key); Bau, 1929a: 4, 1929b: 544 [*Cuterebra (Orthocuterebra)*]; Aldrich, 1931: 117; Bau, 1931b: 206 [*Cuterebra (Pseudobogeria)*]; Townsend, 1931: 79 (as syn. of *princeps* Austen, 1895); Curran, 1942: 79 (in key); Dalmat, 1943a: 182 (as syn. of *princeps* Austen, 1895); Chilcott, 1965: 1110 (as syn. of *princeps* Austen, 1895); J.H. Guimarães, 1967: 3 (cat.; as syn. of *princeps* Austen, 1895); Baird, 1983: 214 (biol.); Sabrosky, 1986: 114, map 5, figs. 28, 34; Pfaffenberger & Valencia, 1988: 842; Colwell, Baird, Lee & Milton, 1999: 803 (SEM and morphometrics of egg).
- sp.* Townsend, 1892: 291.
- fontinella* Clark of Townsend, 1893: 317, 1895a: 64, 1895b: 618, and Honess & Winter, 1956: 239, fig. 115, misident.
- sp.* Austen, 1895a: 384, 1895b: 152 (*Cutiterebra*).
- buccata* Fabricius of Baker, 1904: 35, misident.
- albifrons* Swenk, 1905: 182. Type-locality: U. S. A., Wyoming, Hecla. Ref. – Bau, 1929a: 4, 1929b: 77 [*Cuterebra (Orthocuterebra)*], 1931b: 206 [*Cuterebra (Pseudobogeria)*]; J.H. Guimarães, 1967: 3 (cat.; as syn. of *princeps* Austen, 1895).
- subbuccata* Bau, 1929c: 305, figs. 4, 6. Type-locality: U. S. A., Nevada, Carson City County (as Ormsby County). Ref. – Bau, 1931b: 206 [*Cuterebra (Pseudobogeria)*].
- maculosa** Knab, 1914: 187. Lectotype-locality: Panama, Canal Zone. Distr. – Guatemala, Panama. Refs. – Townsend, 1917: 24 (in key), 1919: 592; Bau, 1931: 205 [*Cuterebra (Cuterebra)*]; Fairchild, 1943: 577; J.H. Guimarães, 1967: 3 (cat.); Sabrosky, 1986: 95.
- postica** Sabrosky, 1986: 140. Type-locality: Mexico, Morelos. Distr. – Mexico (Morelos, Oaxaca).
- praegrandis** Austen, 1933: 706, pl. 1, figs. 6-7. Type-locality: Peru, Loreto, Iquitos. Distr. – Peru (Loreto), Brazil (Pará). Refs. – J. H. Guimarães, 1967: 3 (cat.), 1984: 25 (*Cuterebrella*); J. H. Guimarães & Papavero, 1999: 203, figs. 98, 110, 118.
- princeps** (Austen), 1895: 105, pl. 13, fig. 5 (*Bogeria*) (also puparium described). Type-locality: Mexico, Sonora, Médano Blanco, Santa Bárbara Bay, Gulf of California. Distr. – U. S. A. (Arizona), Mexico (Sonora). Refs. – Brauer, 1895: 591 (*Bogeria*); Coquillett, 1910: 515 (*Bogeria*); Townsend, 1917: 24 (in key), 25 (*Bogeria*); Bau, 1931b: 198, 206 (*Bogeria*); Townsend, 1931: 79 (*Bogeria*), 1938: 169 (*Bogeria*); Curran, 1942: 79 (n. comb.); J.H. Guimarães, 1967: 3 (cat.); Cole, 1970: 586, 587; Sabrosky, 1986: 105, map 4, fig. 31.
- sabroskyi** (J. H. Guimarães), 1984: 26, fig. 5 (*Cuterebrella*). Type-locality: Brazil, São Paulo, São Paulo (Pico do Jaraguá). Distr. – Brazil (São Paulo). Ref. – J. H. Guimarães & Papavero, 1999: 204, fig. 99.
- sp. near albipilosa** Sabrosky, 1986: 179 (U. S. A.: Texas; Mexico, Coahuila, San Luís Potosí).
- terrizona** Walker, 1849: 683. Type-locality: “Guatemala”. Distr. – Central and southern Mexico (Méjico, Nayarit, Oaxaca, Chiapas) south to Guatemala and El Salvador. Refs. – Brauer, 1863b: 230 (in key; as syn. of *americana* Fabricius, 1775); Wulp, 1888: 1 (as syn. of *americana* Fabricius, 1775); Austen, 1895b: 147 (*Cutiterebra*; as syn. of *atrox* Clark, 1848); Swenk, 1905: 182 (in key; as syn. of *atrox* Clark, 1848); Bau, 1928: 51 (as syn. of *atrox* Clark, 1848), 1929a: 3 (in key; as syn. of *atrox* Clark, 1848), 1931b: 207 [*Cuterebra (Atrypoderma)*, as syn. of *atrox* Clark, 1848]; Austen, 1933: 702 (as syn. of *atrox* Clark, 1848); J.H. Guimarães, 1967: 4 (cat.); Sabrosky, 1986: 152, map 11.
- approximata* Walker of Wulp, 1888: 2, misident., part (only his Guatemala record).

Genus **Dermatobia** Brauer, 1861

Dermatobia Brauer, 1861: 783. Type-species, *Cuterebra cyaniventris* Macquart, 1843 (sub. des., J.H. Guimarães & Papavero, 1966: 228) = *hominis* (Linnaeus Jr., 1781).

hominis (Linnaeus Jr.) in Pallas, 1781: 158 (*Oestrus*). Type-locality: “Peru”. Distr. – U. S. A. to Argentina, but not in Chile. Refs. – Gmelin, 1788: 2811 (*Oestrus*); Brauer, 1861: 783; Bau, 1922 (*Oestrus*); J. H. Guimarães, 1966 (aggregation habit of males); J.H. Guimarães, 1967: 5 (cat.); J.H. Guimarães & Papavero, 1999: 261, figs. 2, 76, 107, 109, 128 (history of studies, common names, characterization and distribution, list of vectors, pathogeny, diagnostic, treatment and control, bibliography).

humanus Howship, 1833: 181 (larva) (*Oestrus*) Type-locality: Colombia, Tolima, Mariquita.

cyaniventris Macquart, 1843: 23 (*Cuterebra*). Type-locality: “Brazil”.

noxialis Goudot, 1845: 229, pl. 4bis, figs. 1-5 (*Cuterebra*). Type-locality: Colombia, Caldas, Marmato.

Genus **Metacuterebra** Bau, 1929

Cuterebra, subg. **Metacuterebra** Bau, 1929b: 544. Type-species, *Cuterebra apicalis* Guérin-Ménéville, 1835 (sub. des., Bau, 1931b: 208).

Alouattamyia Townsend, 1931: 319. Type-species, *Cuterebra baeri* Greene, 1925 (orig. des.).

Oricuterebra Bau, 1931b: 210. Type-species, *Cuterebra sarcophagooides* Lutz (orig. des.).

Baucuterebra L. R. Guimarães & Carrera, 1941: 2. Type-species, *Cuterebra schroederi* Enderlein (sub. des., Townsend, 1942 (Addenda & Corrigenda): 333).

Montemyia Dalmat, 1942a: 25. Type-species, *Montemyia bureni* Dalmat, 1942 (orig. des.).

Refs. – J. H. Guimarães, 1971 (hosts); J. H. Guimarães & Papavero, 1999: 211 (key to spp.).

almeidai (L. R. Guimarães & Carrera), 1941: 2, figs. 1-3 (*Baucuterebra*). Type-locality: Brazil, Amazonas, Cachoeira Taurino. Distr. – Brazil (Amazonas, Pará), Argentina. Refs. – J. H. Guimarães, 1967: 2 (cat.); J. H. Guimarães & Papavero, 1999: 213, fig. 90 (this species is probably a variation of *Metacuterebra funebris* (Austen). However, as only a few specimens of *almeidai* are available to evaluate the variation of the chromatic characters of pleural pilosity, we maintain it as a valid species); Pinto & Claps, 2005: 572 (Argentina record).

apicalis (Guérin-Ménéville), 1835: pl. 101, fig. 1 (*Cuterebra*). Type-locality: “America”. Distr. – South Mexico to Argentina (Mendoza). Refs. – Brauer, 1863b: 237, 239 (*Cuterebra*); Wulp, 1888: 2 (*Cuterebra*); Bau, 1906: 25, pl. 2, fig. 17 (*Cuterebra*); Lutz, 1917: 47 (*Cuterebra*); Bau, 1929a: 6, 1929b: 544 (*Cuterebra*), 1931b: 208 (*Cuterebra*); Fonseca, 1938: 314 (*Cuterebra*), 1940: 665; Henriksen, 1942: 284 (*Cuterebra*; hosts, larva); Forattini & Lenko, 1959: 155 (*Cuterebra*; descr. of pupa); d’Andretta Jr. & Jardim, 1954: 67 (*Cuterebra*); Del Ponte, 1959: 755 (*Cuterebra*); J. H. Guimarães, 1967: 2 (cat.), 1971: 90 (hosts) (*Cuterebra*); Led, Colacelli, Boero & Colombo, 1976: 35 (in *Rattus norvegicus*); Mello, 1978: 329 (*Cuterebra*; biol.); Leite, Guimarães & Lima, 1986: 89 (ESM study of egg); Leite & Williams, 1988a: 485 (life cycle), 1989: 189 (egg and first instar larva); Bossi & Bergallo, 1992: 142 (in *Oryzomys nitidus* (Rodentia: Cricetidae) and *Metachirus nudicaudatus* (Marsupialia: Didelphidae)); Vieira, 1993: 792 (hosts in “cerrado” [savanna] from Central Brazil), J.H. Guimarães & Papavero, 1999: 213, figs. 100, 105, 117; Pujol-Luz, Mendonça & Henriques, 2004: 669 (*Gracilinanus agilis* (Marsupialia, Didelphidae) as host, Brasília, D. F.); Pinto & Claps, 2005: 572 (Argentina, in *Oligoryzomys longicaudatus* and *Akodon caenosus* (Rodentia: Muridae)).

analis Macquart, 1843: 22, pl. 2, fig. 5 (*Cuterebra*). Type-locality: “Brazil”. Refs. – Joly, 1846: 278, fig. 7; Brauer, 1863b: 237, pl. 4, fig. 1, pl. 6, fig. 8; Austen, 1895a: 386 (*Cutiterebra*).

cometes Shannon & Del Ponte, 1926: 42 (*Cuterebra*). Type-locality: Argentina, Tucumán, Quebrada de Lules. Refs. – Del Ponte, 1939: 527; J. H. Guimarães, 1967: 2 (cat.); J. H. Guimarães & Papavero, 1999: 213 (syn.).

pusilla Bau, 1931b: 212, pl. 17, fig. 2, pl. 2, fig. 16 (*Cuterebra*). Type-locality: Bolivia, Mapiri, Sarampiuni. Refs. – Ref. – J.H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J.H. Guimarães & Papavero, 1999: 213 (syn.).

baeri (Greene), 1925: 160 (*Cuterebra*). Type-locality: Guyana, Bartica, Kartabo. Distr. – Costa Rica, Panama, Guyana, Peru, Brazil (Amazonas, Pará). Refs. – Shannon & Greene, 1926: 286, pls. 152-153 (*Cuterebra*); Townsend, 1931: 319 (*Alouattamyia*); Carpenter, 1934; Dunn, 1934; Zeledón, Jiménez & Brenes, 1957: 129 (*Cuterebra*; larva); J. H. Guimarães, 1967: 1 (cat., *Alouattamyia*), 1971 (*Alouattamyia*); J. H. Guimarães & Coimbra, 1982: 35; Fraiha Neto, Chaves, Borges & Freitas, 1984: 63 (*Alouattamyia*); Dudley & Milton, 1990 (*Alouattamyia*); Baron, Colwell &

- Milton, 1996: 946 (*Alouattamyia*); Milton, 1996: 39 (*Aloattamyia*); Colwell & Milton, 1998: 674 (*Alouattamyia*; development in *Alouatta palliata*); Stuart, Pendergast, Rumfelt, Pierberg, Greenspan, Glander & Clarke, 1998 (*Alouattamyia*); J. H. Guimarães & Papavero, 1999: 222, figs. 87, 111, 123 (n. comb.); Colwell, Baird, Lee & Milton, 1999: 803 (*Alouattamyia*; SEM and morphometrics of egg).
- bureni** (Dalmat), 1942a: 25, pl. 2 (*Montemyia*). Type-locality: Costa Rica, Monte Redondo. Distr. – Costa Rica. Ref. – J. H. Guimarães, 1967: 5 (cat.).
- cayennensis** (Macquart), 1843: 23 (*Cuterebra*). Type-locality: French Guiana, Cayenne. Distr. – Nicaragua, Panama, Surinam, French Guiana, Brazil (Pará). Refs. – Joly, 1846: 289 (*Cuterebra*); Brauer, 1863b: 240 (*Cuterebra*); J. H. Guimarães, 1967: 2 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 224, figs. 86, 93, 112.
- pygmaea** Bau, 1931b: 251, pl. 5, fig. 5 (*Cuterebra*). Type-locality: Brazil, Pará, ilha de Marajó. Refs. – J. H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 224 (syn.).
- levida** Austen, 1933: 707, pl. 1, fig. 8 (*Cuterebra*). Type-locality: “Nicaragua”. Refs. – J. H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 224 (syn.).
- detrudatrix** (Clark), 1848: fig. 4 (*Cuterebra*, as *detrudator*). Type-locality: “South America”. Distr. – Panama, Trinidad, Surinam, Ecuador, Peru, Brazil (Amazonas, Pará, Espírito Santo, São Paulo, Paraná). Refs. – Brauer, 1863b: 49 (*Cuterebra*); Bau, 1931b: 233 (*Cuterebra*); J. H. Guimarães, 1967: 2 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 225, figs. 104, 114.
- nigricincta** Austen, 1895a: 388, pl. 13, figs. 4, 4a (*Cutiterebra*). Type-locality: Brazil, Pará. Refs. – Bau, 1931b: 233 (*Cuterebra*); Austen, 1895b: 149 (*Cutiterebra*; syn.), 1933: 710; J. H. Guimarães, 1967: 3 (cat.; *Cuterebra*).
- schroederi** Enderlein, 1909: 202 (*Cutiterebra*). Type-locality: Peru, Chanchamayo. Refs. – Bau, 1929c: 307, fig. 5; J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 225 (syn.).
- nigricans** Lutz, 1917: 99 (1918: 123), pl. 27, fig. 2, pl. 29, fig. 2 (*Cuterebra*). Type-locality: Brazil, São Paulo, Porto Martins. Ref. – J. H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 225 (syn.).
- conflans** Bau, 1929c: 303, figs. 1-2 (*Cuterebra*; as *conflans conflans*). Type-locality: Peru, Junín, Chanchamayo. Refs. – Bau, 1931b: 225; J. H. Guimarães, 1967: 2 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 225 (syn.).
- obidosiana** Bau, 1929c: 305 (*Cuterebra*; as *conflans* spp.). Type-locality: Brazil, Pará, Óbidos. Refs. – Bau, 1931b: 233 (*Cuterebra*); J. H. Guimarães, 1967: 2 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 225 (syn.).
- dorsalis** Bau, 1929c: 305, fig. 3 (1930: 83) (*Cuterebra*). Type-locality: “South America”. Refs. – J. H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 225 (syn.).
- pulchra** Bau, 1930: 85, fig. 4 (*Cuterebra*). Type-locality: “South America”. Refs. – J. H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 225 (syn.).
- versicolora** Bau, 1930: 85, fig. 4 (*Cuterebra*, as *versicolora versicolora*). Type-locality: “Surinam”. Refs. – J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 225 (syn.).
- bicincta** Bau, 1931b: 221, pl. 1, fig. 6, pl. 2, fig. 17 (*Cuterebra*). Type-locality: “Surinam”. Refs. – J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*, as *versicolora* ssp.); J. H. Guimarães & Papavero, 1999: 225 (syn.).
- worontzowi** L. R. Guimarães & Carrera, 1941: 6, figs. 7-9 (*Cuterebra*). Type-locality: Brazil, Amazonas, rio Paruari. Refs. – J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 225 (syn.).
- funebris** (Austen), 1895a: 378, pl. 13, figs. 1, 16 (*Cutiterebra*). Type-locality: “Trinidad”. Distr. – Trinidad, Brazil (Goiás). Refs. – Bau, 1931b: 230 (*Cuterebra*); Austen, 1933: 704 (*Cuterebra*); J. H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 227.
- ribeiroi** Artigas & d’Andretta Jr., 1951: 60, figs. 1-3 (*Alouattamyia*). Type-locality: Brazil, Goiás, Aruanã, rio Vermelho. Ref. – J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*).
- infulata** (Lutz), 1917: 99, pl. 27, fig. 4, pl. 29, fig. 4 (*Cuterebra*). Type-locality: Brazil, Rio de Janeiro, Petrópolis. Distr. – Brazil (Pernambuco, Rio de Janeiro, São Paulo, Rio Grande do Sul). Refs. – J. H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 227, figs. 92, 121.
- bauii** Fonseca, 1940: 667, figs. 1-3 (*Oicuterebra*). Type-locality: Brazil, São Paulo. Refs. – J. H. Guimarães, 1967: 2 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 227 (syn.).
- rubiginosa** Bau, 1931b: 214, pl. 1, fig. 9, pl. 2, fig. 15 (*Cuterebra*). Type-locality: Brazil, Rio Grande do Sul, Santa Cruz. Refs. – J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 227 (syn.).
- megastoma** (Brauer), 1863: 247, pl. 4, fig. 5, pl. 6, fig. 12 (*Cuterebra*). Type-locality: “South America”. Distr. – Brazil (Minas Gerais, São Paulo, Rio de Janeiro, Santa Catarina, Rio Frande do Sul). Refs. – J. H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 228, figs. 95, 119.
- sarcophagoides** Lutz, 1917: 99 (1918: 123), pl. 27, fig. 5, pl. 29, fig. 5. Type-locality: Brazil, São Paulo, Jacutinga. Refs. – J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 228 (syn.).
- patagona** (Guérin-Ménéville), 1844: 548 (*Cuterebra*). Type-locality: Argentina, Patagonia. Distr. – Argentina (Misiones, Chubut, La Pampa, Santa Cruz). Refs. – Del Ponte, 1939: 527 (*Cuterebra*); J. H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 229, fig. 88.

- pessoai** (L. R. Guimarães & Carrera), 1941: 4, figs. 4, 6 (*Cuterebra*). Type-locality: Brazil, Amazonas, rio Negro. Distr. –Trinidad, Brazil (Amazonas). Refs. – J.H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 230, fig. 91.
- rufiventris** (Macquart), 1843: 21, figs. 4, 4b (*Cuterebra*). Type-locality: Brazil, Pará [*i.e.*, Belém]. Distr. – Ecuador, Peru, Brazil (Amazonas, Pará, Mato Grosso do Sul, Rio de Janeiro, Santa Catarina). Refs. – Brauer, 1863b: 245 *Cuterebra*; Austen, 1895a: 387, pl. 13, figs. 3, 3a (*Cutiterebra*); Bau, 1906: 26 (*Cuterebra*); Lutz, 1917: 89 (*Cuterebra*); Bau, 1931b: 225 (*Cuterebra*); Austen, 1933: 705 (*Cuterebra*); J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 231, fig. 120.
- schmalzi** Lutz, 1917: 100 (1918: 124), pl. 27, fig. 3, pl. 24, fig. 3 (*Cuterebra*). Type-locality: Brazil, Santa Catarina, Joinville. Refs. – Bau, 1929a: 7 (*Cuterebra*); J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*); Leite, Falquete & Guimarães, 1985: 835 (*Cuterebra*; host); J. H. Guimarães & Papavero, 1999: 231 (syn.).
- elegans** Bau, 1931b: 227 (*Cuterebra*, as *rufiventris* ssp.). Type-locality: Peru, Junín, Chanchamayo. Refs. – J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*, as *rufiventris* ssp.); J. H. Guimarães & Papavero, 1999: 231 (syn.).
- simulans** (Austen), 1933: 709 (*Cuterebra*). Type-locality: “Peru”. Distr. – Surinam, Peru, Brazil (Pará). Refs. – J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 2302, fig. 89.
- townsendi** (Fonseca), 1941: 483, fig. 1 (*Oricuterebra*). Type-locality: Brazil, São Paulo, São Paulo. Distr. – Brazil (São Paulo). Refs. – J. H. Guimarães, 1967: 4 (cat.; *Cuterebra*); J. H. Guimarães & Papavero, 1999: 233, figs. 94, 106, 115.

Unrecognized or unidentified species of *Cuterebra* and *Metacuterebra*

- ephippium** Latreille, 1818: 271 (*Cuterebra*). Type-locality: French Guiana, Cayenne. Refs. – J.H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J.H. Guimarães & Papavero, 1999: 234.
- flaviventris** Bau, 1931b: 219, pl. 1, fig. 10 (*Metacuterebra*). Type-locality: “Mexico”. Ref. – J.H. Guimarães, 1967: 3 (cat.; *Cuterebra*).
- lutzi** Bau, 1930: 81 (*Cuterebra*). Type-locality: “Peru” and “Brazil”. Refs. – J.H. Guimarães, 1967: 3 (cat.; *Cuterebra*); J.H. Guimarães & Papavero, 1999: 234.
- ornata** Bau, 1928a: 50 (*Cuterebra*). Type-locality: “Costa Rica”. Ref. – J.H. Guimarães, 1967: 3 (cat.; *Cuterebra*).
- semilutea** Bau, 1929a: 1, figs. 1-2 (*Cuterebra*). Type-locality: “Brazil”. Ref. – J.H. Guimarães, 1967: 4 (cat.; *Cuterebra*); J.H. Guimarães & Papavero, 1999: 234.
- sp.** Adler, Davis & Carvajal, 2003: 693 (*Cuterebra*). [Panama, in *Proechimys semispinosus* (Rodentia, Echimyidae)].
- sp.** Led & Brandelli, 1978: 29 (*Cuterebra*) [Chile, in *Chinchilla lanigera* (Rodentia, Chinchillidae)].
- sp.** Manrique-Saide, Hernandez-Betancourt & Quintero, 2000: 487 (*Cuterebra*) [Mexico, Yucatán; in *Ototylomys phyllotis* (Rodentia: Muridae)].

Genus *Pseudogametes* Bischoff, 1900

- Pseudogametes** Bischoff, 1900a: 131 (1900b: 490). Type-species, *Pseudogametes hermanni* Bischoff, 1900 (orig. des.).
Xylocopodes Townsend, 1927: 207. Type-species, *Musca semiatra* Wiedemann, 1830 (orig. des.). Ref. – J. H. Guimarães, 1967: 5 (cat.; syn.).
- Ref. – J. H. Guimarães & Papavero, 1999: 205 (key to spp.).

- hermanni** Bischoff, 1900a: 131 (1900b: 490, figs. 1-3). Type-locality: Brazil, Minas Gerais. Refs. – Lutz, 1917: 102 (1918: 126), pl. 28, fig. 7; J. H. Guimarães, 1967: 5 (cat.); J. H. Guimarães & Papavero, 1999: 206, figs. 102, 108.
- semiater** (Wiedemann), 1830: 421 (*Musca*). Type-locality: “Brazil”. Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo). Refs. – Lutz, 1917: 102 (1918: 126); Townsend, 1927: 207 (as *Xylocopodes semiatra*); J. H. Guimarães, 1967: 5 (cat.); J. H. Guimarães & Papavero, 1999: 205, fig. 122.

Genus *Rogenhofera* Brauer, 1863

- Rogenhofera** Brauer, 1863a: 325 (1863b: 215, 217). Type-species, *Rogenhofera trigonophora* Brauer, 1862 (mon.).
Rogenhoferia, error or emend.

- Guerinioestrus* Townsend, 1931: 319. Type-species, *Cephenemyia grandis* Guérin-Ménéville,
Refs. – Scaglia & Velásquez, 1982; J. H. Guimarães & Papavero, 1999: 207 (key to spp.).
- dasypoda** Brauer, 1896: 272. Type-locality: Brazil, Espírito Santo. Distr. – Brazil (Espírito Santo, Rio de Janeiro, São Paulo).
Refs. – Lutz, 1917: 100 (1918: 124), pl. 28, fig. 8; Bau, 1928b: 300; J. H. Guimarães, 1967: 5 (cat.), 1990: 256; J. H. Guimarães & Papavero, 1999: 207, figs. 97, 103, 106.
- gilvopilosa** Bau, 1932: 11. Type-locality: Peru, Arequipa, Puno. Distr. – Peru (Arequipa). Refs. – J. H. Guimarães, 1967: 5 (cat.); J. H. Guimarães & Papavero, 1999: 208.
- grandis** (Guérin-Ménéville), 1844: 548 (*Cephenemyia*). Type-locality: Argentina, Patagonia. Distr. – Argentina (Córdoba, Buenos Aires, Patagonia). Refs. – Brauer, 1863b: 213; Bau, 1906: 24; Townsend, 1931: 318; Bau, 1932: 11; Del Ponte, 1939: 528; J. H. Guimarães, 1967: 5 (cat.); J. H. Guimarães & Papavero, 1999: 108, fig. 113.
- bonaerensis** Del Ponte, 1939: 529 (*Guerinioestrus*). Type-locality: Argentina, Buenos Aires. Refs. – J. H. Guimarães, 1967: 4 (cat.); Brigada, Tripole & Zuleta, 1992: 646 (in *Akodon molinae* (Rodentia: Cricetidae), Argentina); J. H. Guimarães & Papavero, 1999: 208 (syn.); Zuleta & Vignau, 1990: 11 (biol.).
- daguerrei** Del Ponte, 1939: 530. Type-locality: “Argentina”. Refs. – Del Ponte, 1959: 575; J. H. Guimarães, 1967: 5 (cat.); J. H. Guimarães & Papavero, 1999: 208 (syn.).
- lopesi** J. H. Guimarães, 1990: 256. Type-locality: Brazil, Goiás, Parque Nacional das Emas. Distr. – Brazil (Goiás). Ref. – J. H. Guimarães & Papavero, 1999: 209.
- trigonophora** Brauer, 1863a: 326 (1863b: 217, pl. 6, fig. 14, pl. 10, fig. 5). Type-locality: Brazil, Bahia. Distr. – Brazil (Bahia, Rio de Janeiro, São Paulo). Refs. – Lutz, 1917: 101 (1918: 125); J. H. Guimarães, 1967: 5 (cat.); J. H. Guimarães & Papavero, 1999: 210.
- trigonocephala** Lutz, 1917: 96, error.

Nomina nuda

- jorgensenii** Del Ponte, 1959: 575 (*Metacuterebra*).
vianai Del Ponte, 1959: 575 (*Guerinioestrus*).

References

- Adler, G. H., S. L. Davis & A. Carvajal, 2003. Bots (Diptera: Oestridae) infesting a Neotropical forest rodent, *Proechimys semispinosus* (Rodentia: Echimyidae), in Panama. *J. Parasitol.* 89 (4): 693-697.
- Agassiz, L., 1846. *Nomenclator zoologicus, nomina systematica generum animalium tam viventium quam fossilium, secundum ordinem alphabeticum disposita, adjectis auctoribus, libris in quibus reperiuntur, anno editionis, etymologia et familias ad quas pertinent in variis classibus. Index.* Soloduri [= Solothurn].
- Alcock, J. & D. J. Kemp, 2004. Long-term stability in the mating system of the bot fly *Cuterebra austeni* (Cuterebridae). *J. Insect Behavior* 17 (3): 273-280.
- Aldrich, J. M., 1905. A catalogue of North American Diptera. *Smithson. Misc. Collns* 46 [2 (= publ. 1444)]: 1-680.
- Aldrich, J. M., 1931. Notes on Diptera No. 5. *Proc. ent. Soc. Washington* 33: 116-121.
- d'Andretta Jr., C. & J. L. Jardim, 1954. Contribuição ao conhecimento do ciclo biológico de *Metacuterebra apicalis* (Guérin 1829/38 [sic]) (Diptera, Cuterebridae). *Anais Fac. Farm. Odontol. Univ. São Paulo* 12: 67-73.
- Artigas, P. T. & C. d'Andretta Jr., 1951. *Alouattamyia ribeiroi*, nova espécie de Cuterebridae (Diptera) do Brasil. *Archos Mus. nac.*, Rio de Janeiro 42: 59-62, 3 pls.
- Austen, E. E., 1895a. On the specimens of the genus *Cutiterebra* and its allies (family Oestridae) in the collection of the British Museum, with descriptions of a new genus and three new species. *Ann. Mag. nat. Hist.* 6 (13): 377-396, 1 pl.
- Austen, E. E., 1895b. Further notes on *Cutiterebra*. On the identity of certain species described by the late Bracy Clark. *Ann. Mag. nat. Hist.* (6) 16: 147-155.
- Austen, E. E., 1933. New and little known species of *Cuterebra* Clark and *Bogeria* Austen (Diptera: Oestridae). *Proc. zool. Soc. London* 3: 699-713, 1 pl.
- Baird, C. R., 1983. Biology of *Cuterebra lepusculi* Townsend (Diptera: Cuterebridae) in cottontail rabbits in Idaho. *J. Wildlife Dis.* 19 (3): 214-218.
- Baird, C. R., 1997. Bionomics of *Cuterebra austeni* (Diptera: Cuterebridae) and its association with *Neotoma albigenula* (Rodentia: Cricetidae) in the southwestern United States. *J. med. Ent.* 34 (6): 690-695.
- Baird, C. R., J. K. Podgore & C. W. Sabrosky, 1982. *Cuterebra* myiasis in humans: Six new cases reports from the United States with a summary of known cases (Diptera: Cuterebridae). *J. med. Ent.* 19: 263-267.
- Baird, J. K., C. R. Baird & C. W. Sabrosky, 1989. North American cuterebrid myiasis. Report of seventeen new infections of human beings and review of the disease. *J. American Acad. Dermatol.* 21: 763-772.
- Baker, C. F., 1904. Reports on Californian and Nevadan Diptera. I. *Invertebrata pacifica* 1: 17-40.
- Baron, R. W., D. D. Colwell & K. Milton, 1996. Antibody immunoglobulin G (IgG) response to *Alouattamyia baeri* (Diptera, Cuterebridae) parasitism of howler monkeys, *Alouatta palliata*, in Panama. *J. econ. Ent.* 33: 946-951.
- Barrett, G. W. & G. A. Felhammer, 2008. Ectoparasites, bots, and vector-borne diseases associated with the golden mouse, pp. 167-189, in their *The golden mouse. Ecology and conservation*. Springer, New York.
- Bau, A., 1906. Diptera, Fam. Muscaridae, subfam. Oestrinae, in Wytsman, P., *Gen. Ins.* 43: 1-31, 2 pls.
- Bau, A., 1922. Ueber die technisch-biologische Bedeutung der Oestriden, das verirrte Vorkommen ihrer Larven, sowie über *Oestrus hominis*, *Hypoderma equi* und *alcis*. *Zeitschr. techn. Biol.* 10 (3-4): 177-193.
- Bau, A., 1928a. *Cuterebra ornata*, spec. nov. *Konowia* 7: 50-51.
- Bau, A., 1928b. Ueber *Aulacocephala* und *Rogenhofera*. *Konowia* 7: 298-303.
- Bau, A., 1929a. *Cuterebra semilutea*, spec. nov. (Diptera), sowie Schlüssel zur Bestimmung der *Cuterebra*-Arten. *Senckenbergiana* 11 (1-2): 1-7, figs. 1-2.
- Bau, A., 1929b. Versuch einer Teilung der Gattung *Cuterebra* (Diptera, olim Oestridae) in vier Untergattungen. *ZentBlatt Bakteriol. ParasitKde Infektionkr.* 77 (25-26): 542-544, 1 fig.
- Bau, A., 1929c. *Cuterebra conflans* und *subbuccata*, spec. nov., sowie Bemerkung über *C. schroederi* Enderlein (Diptera, olim Oestridae). *Stettin. ent. Ztg* 90: 303-307, 1 pl.
- Bau, A., 1930. Vier neue *Cuterebra*-Arten aus Südamerika (Diptera, olim Oestridae). *Konowia* 9 (1): 81-89, pl. 1. figs. 1-6.
- Bau, A., 1931a. *Cuterebra (Atrypoderma) enderleini* spec. nov. (Dipt.). *Mitt. deutsche ent. Ges.* 2: 20-21.
- Bau, A., 1931b. Ueber das Genus *Cuterebra* Clark (Diptera, Oestridae). Einteilung derselben in sechs Untergattungen. Beschreibung neuer Species und Aufstellung einer Bestimmungstabelle der mittel- und südamerikanischen Arten. *Konowia* 10: 197-240, 2 figs., 2 pls.
- Bau, A., 1932. *Rogenhofera gilvopilosa*, spec. n. und *R. grandis* (Guérin) (Dipt. Cuterebr.). *Mitt. deutsche ent. Ges.* 3: 11-15.
- Beamer, R. H., 1950. An observation on the egg-laying of *Cuterebra buccata* Fabr. in nature. *J. Kansas ent. Soc.* 23: 16.
- Benchimol, J.L. & M.R. Sá. 2006. *Adolpho Lutz. Obra completa.* 4 (2). *Entomologia*. 1053 pp. Editora Fiocruz, Rio de Janeiro.
- Bequaert, J., 1928. Cuterebridae, p. 805, in Johannsen, O. A., Order Diptera, in Leonard, A list of the insects of New York. *Cornell Univ. Agric. Expt Sta. Mem.* 101: 1-1121.
- Bequaert, J., 1945. Cutaneous myiasis due to *Cuterebra* in Massachusetts. *Psyche* 52: 175-176.
- Berg, C., 1876. Naturgeschichte der *Rogenhofera grandis*, einer Fliege aus der Familie der Oestriden. *Stettin. ent. Ztg* 37: 268-272.

- Bischoff, J., 1900a. Vorläufige Charakteristik einiger neuen Gattungen von Muscarien. *Anz. k. Akad. Wiss. Wien* 37: 131.
- Bischoff, J., 1900b. Einige neue Gattungen von Muscarien. *SitzBer. k. Akad. Wiss. Wien, math.-naturw. Cl.* 109: 490-497, 5 figs.
- Berg, C., 1881. Entomologisches aus dem Indianergebiet der Pampa. *Stettin. ent. Ztg* 42 (1): 36-72.
- Blanchard, C. E., 1840. Histoire naturelle des insectes. Orthoptères, névroptères, hémiptères, hyménoptères, lépidoptères et diptères. Vol. 3, 672 pp., 67 pls., in Laporte, F. L. de, *Histoire naturelle des animaux articulés*. Paris.
- Blanchard, C. E., 1845. *Histoire des insectes* 2: 524 pp., pls. 11-20. F. Didot Frères, Paris.
- Bossi, D. E. P. & H. G. Bergallo, 1992. Parasitism by cuterebrid botflies (*Metacuterebra apicalis*) in *Oryzomys nitidus* (Rodentia: Cricetidae) and *Metachirus nudicaudatus* (Marsupialia: Didelphidae) in a southeastern Brazilian rainforest. *J. Parasitol.* 78 (1): 142-145.
- Brauer, F. M., 1861. Ueber die Larven der Gattung *Cuterebra*. *Clk. Verh. zool.-bot. Ges. Wien* 10 (1860): 777-786.
- Brauer, F. M., 1863a. *Rogenhofera*, eine neue Gattung aus der Familie der Oestriden. *Verh. zool.-bot. Ges. Wien* 13: 325-326.
- Brauer, F., 1863. *Monographie der Oestriden*, vi + 292 pp., 10 pls. C. Ueberreuter, Wien. [Also publ. in *Verh. zool.-bot. Ges. Wien* 13 (Beih.): vi + 291 pp.].
- Brauer, F., 1887. Nachträge zur Monographie der Oestriden. II. Zur Charakteristik und Verwandtschaft der Oestriden Gruppen im Larven- und volkommenen Zustande. *Wien. ent. Ztg* 6: 4-16.
- Brauer, F. M., 1895. Bemerkungen zu einigen neuen Gattungen der Muscarien und Deutung einiger Original-Exemplare. *SitzBer. Akad. Wiss. Wien* 104 (Abt. I): 582-604.
- Brauer, F. M., 1896. Beiträge zur Kenntniss aussereuropäische Oestriden. *Denkschr. k. Akad. Wiss. Wien, math.-naturw. Cl.* 64: 260-282, 1 pl.
- Brigada, A. M., E. S. Tripole & G. A. Zuleta, 1992. Cuterebrid parasitism (*Rogenhofera bonaerensis*) on the shrubland mouse (*Akodon molinae*) in Argentina. *J. Wildl. Dis.* 28 (4): 646-650.
- Brimley, C. S., 1938. *The insects of North Carolina. Being a list of the insects of North Carolina and their close relatives*, 560 pp. North Carolina Department of Agriculture, Raleigh.
- Bromley, S. W., 1937. New and little-known Utah Diptera with notes on the taxonomy of the Diptera. *Proc. Utah Acad. Sci.* 14: 99-109.
- Burt, W. H., 1940. Territorial behavior and populations of some small mammals in southern Michigan. *Misc. Publs Mus. Zool. Univ. Michigan* 45: 1-58, 2 pls.
- Cameron, A. E., 1926. The occurrence of *Cuterebra* (Diptera, Oestridae) in western Canada. *Parasitology* 18: 430-435.
- Carpenter, C. R., 1934. A field study of the behavior and social relations of howling monkeys. *Comp. Psychol.* Mon. 10: 1-168.
- Catts, E. P., Jr., 1982. Biology of New World bot flies: Cuterebridae. *Annu. Rev. Ent.* 27: 313-338, 2 figs., 1 table.
- Chilcott, J. F., 1965. Family Cuterebridae, pp. 1109-1110, in Stone, A., C. W. Sabrosky, W. W. Wirth & J. R. Coulson, eds., *A catalog of the Diptera of America North of Mexico*. [U. S. Dept. Agric., Agric. Res. Serv.] *Handbook* 276: 1696 pp. Washington, D. C.
- Clark, B., 1815. *An essay on the bots of horses and other animals*, 72 pp., 2 pls. London.
- Clark, B., 1827. On the insect called Oistros by the ancients. To which is added, a description of a new species of *Cuterebra*. *Trans. Linn. Soc. London* 15: 402-411.
- Clark, B., 1843. An appendix or supplement to a treatise on the Oestri and Cuterebrae of various animals. *Trans. Linn. Soc. London* 19: 81-94.
- Clark, B., 1848. Addenda 1848 [To Essay on bots], 1 p., unnumb. London.
- Cogley, T. P., 1991. Warble development by the rodent bot *Cuterebra fontinella* (Diptera: Cuterebridae) in the deer mouse. *Vet. Parasitol.* 38 (4): 275-288.
- Cole, 1970. *The flies of western North America*, xii + 693 pp. Berkeley & Los Angeles ("1969").
- Cole, F. R. & A. L. Lovett, 1921. An annotated list of the Diptera (flies) of Oregon. *Proc. California Acad. Sci.* (4) 11: 197-344.
- Colwell, D. D., C. R. Baird, B. Lee & K. Milton, 1999. Scanning electron microscopy and comparative morphometris of eggs from six bot fly species (Diptera: Oestridae). *J. med. Ent.* 36 (6): 803-810.
- Colwell, D. D. & K. Milton, 1998. Development of *Alouattamyia baeri* (Diptera: Oestridae) from howler monkeys (Primates: Cebidae) on Barro Colorado Island, Panama. *J. med. Ent.* 35 (5): 647-680.
- Coquillett, D. W., 1898. On *Cuterebra emasculator*, with descriptions of several allied species. *Canadian Entomologist* 30: 9-11.
- Coquillett, D. W., 1902. New Diptera from North America. *Proc. U. S. natl Mus.* 25: 83-126.
- Coquillett, D. W., 1910. The type-species of the North American genera of Diptera. *Proc. U. S. natl Mus.* 37 (1719): 499-647.
- Cornet, M., M. Florent, A. Lefebvre, C. Wertheimer, C. Perez-Eid, M. J. Bangs & A. Bouvet, 2003. Tracheopulmonary myiasis caused by a mature third-instar *Cuterebra* larva: Case report and review. *J. clin. Microbiol.* 41 (12): 5810-5812.
- Cramer, M. J. & G. N. Cameron, 2006. Effects of bot fly (*Cuterebra fontinella*) parasitism on a population of white-footed mice (*Peromyscus leucopus*). *J. Mammal.* 87 (6): 1103-1111.
- Cramer, M. J. & G. N. Cameron, 2007. Effects of bot fly, *Cuterebra fontinella*, parasitism on male aggression and female choice in *Peromyscus leucopus*. *Animal Behaviour* 74: 1419-1427.
- Curran, C. H., 1934. *The families and genera of North American Diptera*, 512 pp. New York.
- Curran, C. H., 1942. American Diptera. *Bull. American Mus.*

- nat. Hist.* 80: 51-84.
- Dalmat, H. T., 1942a. New genus and species of Cuterebridae (Diptera) from Costa Rica and other taxonomic notes. *Trans. Amer. ent. Soc.* 68 (1-2): 23-29, pl.
- Dalmat, H. T., 1942b. A new parasitic fly (Cuterebridae) from the northern white-footed mouse. *J. New York ent. Soc.* 50: 45-59.
- Dalmat, H. T., 1942c. A new *Cuterebra* (Diptera: Cuterebridae) from Iowa with notes on certain facial structures. *Amer. Midland Naturalist* 27: 418-421.
- Dalmat, H. T., 1943a. Synonymic notes on some species of *Cuterebra*. *Ent. News* 54 (8): 182-183.
- Dalmat, H. T., 1943b. A contribution to the knowledge of the rodent warble flies (Cuterebridae). *J. Parasitol.* 29: 311-318.
- Davis, W. B., 1944. Notes on Mexican mammals. *J. Mammal.* 25: 370-403.
- Davis, W. T., 1937. [Specimen of the rabbit bot fly, *Cuterebra buccata*]. *Bull. Brooklyn ent. Soc.* 32 (2): 85.
- Del Ponte, E., 1939. Revisión de los "Oestridae" argentinos. *Physis*, Buenos Aires 17: 525-534.
- Del Ponte, E., 1959. Diptera Cuterebridae, p. 575, in Bejarano, J.F.R., E. Del Ponte & R.N. Orfila, *Primeras Jornadas entomoparásiticas argentinas*. Vol. 2. Dirección General de Sanidad de la Secretaría de la Guerra, Buenos Aires.
- Desmarest, E., 1859. Annelés, iv + 312 pp., 40 pls., in Chenu, J. C., *Encyclopédie d'histoire naturelle*. Maresq, Paris.
- Disney, R. H. L., 1969. Notes on rodents warble-flies from British Honduras including the description of a new species of *Cuterebra* Clark (Diptera, Cuterebridae). *Ent. monthly Mag.* 104: 189-197, 3 figs.
- Dixon, J. M., C. H. Winkler & J. H. Nelson, 1969. Ophthalmomyiasis interna caused by *Cuterebra* larva. *Trans. Amer. ophthalmol. Soc.* 67: 110-115.
- Drummond, R. O., R. E. Gingrich & W. J. Gladney, 1970. A new test for detecting animal systemic insecticides. *J. Parasitol.* 56 (4, Sect. 2, pt. 1): 85-86.
- Dudley, R. & K. Milton, 1990. Parasite deterrence and the energetic costs of slapping in howler monkeys, *Alouatta palliata*. *J. Mammal.* 71 (3): 463-465.
- Dunn, L. H., 1934. Entomological investigations in the Chiriquí region of Panama. *Psyche* 41: 166-183.
- Durden, L. A., 1995. Bot fly (*Cuterebra fontinella fontinella*) parasitism of cotton mice (*Peromyscus gossypinus*) on St. Catherine's Island, Georgia. *J. Parasitol.* 81 (5): 787-790.
- Eddy, G. W. & K. S. Emerson, 1940. Notes on Oklahoma botflies. *J. Kansas ent. Soc.* 13: 44-45.
- Enderlein, G., 1909. *Cutiterebra schroederi*, eine neue parasitäre Fliege (Oestridae olim). *Stettin. ent. Ztg* 70 (1): 202-203.
- Everard, C. O. R. & T. H. G. Aitken, 1972. Cuterebrid flies from small mammals in Trinidad. *J. Parasitol.* 58 (1): 189-190.
- Ewing, N. E., 1925. (*Cuterebra* larvae in skin of Panama howler monkey (*Alouatta palliata inconsonans*)). *J. Parasitol.* 12 (2): 107.
- Fabricius, J. C., 1777. *Genera insectorum eorumque characteres naturales secundum numerum, figuram,* situm et proportionem omnium partium oris adiecta mantissa specierum nuper detectarum, 310 pp. Chilonii [=Kiel].
- Fabricius, J. C., 1781. *Species insectorum exhibentes eorum differentias specificas, synonyma, auctorum, loca natalia, metamorphosin* 2: 517 pp. Hamburgi et Kilonii [=Hamburg and Kiel].
- Fabricius, J. C., 1787. *Mantissa insectorum sistens species nuper detectarum* 2: 382 pp. Hafniae [=Copenhagen].
- Fabricius, J. C., 1794. *Entomologia systematica emendata et aucta* 4: 472 pp. Hafniae [=Copenhagen].
- Fabricius, J. C., 1805. *Systema Antliatorum secundum ordines, genera, species*, xiv + 372 + 30 pp. Brunsvigae [=Brunswick].
- Fairchild, G. B., 1943. An annotated list of the bloodsucking insects, ticks and mites known from Panama. *American J. trop. Med. Hyg.* 23: 569-591.
- Fletcher, J. & A. Gibson, 1908. Entomological record, 1907. *Annu. Rept ent. Soc. Ontario* (1907)
- Fonseca, F. O. R. da, 1938. Observação de uma fase do ciclo evolutivo de *Cuterebra apicalis* Guérin (Diptera, Oestridae). *Bolm biol. Clube zool. Brasil* (n. s.) 3 (3-4): 166-167, figs. [Also publ. in *Mems Inst. Butantan*, São Paulo 12 (1938-1939): 195-196, pl., 1939].
- Fonseca, F. O. R. da, 1940. Nova espécie de *Oricuterebra* do Brasil (Dipt., Oestridae). *Revta Ent.*, Rio de Janeiro 11: 662-671, 3 figs.
- Fonseca, F. O. R. da, 1941. Cuterebrineo novo parasito de didelfideo (Diptera, Oestridae). *Revta Ent.*, Rio de Janeiro 12: 480-486, 1 fig., 2 pls.
- Forattini, O. P. & K. Lenko, 1959. Nota biológica sobre *Metacuterebra apicalis* (Guérin, 1823/38 [sic]) (Diptera, Cuterebridae). *Arqos Fac. Hig. Saúde públ. Univ. São Paulo* 13: 155-158, 2 figs.
- Fraiha, H., L. C. L. Chaves, I. C. Borges & R. B. Freitas, 1984. Miases humanas na Amazônia. III. Miase pulmonar por *Alouattamyia baeri* (Shannon & Greene, 1926) (Diptera, Cuterebridae). *Revta Fund. SESP* 29: 63-68.
- Getz, L. L., 1970. Botfly infestations in *Microtus pennsylvanicus* in southern Wisconsin. *American Midland Naturalist* 84 (1): 186-197, 5 figs.
- Gingrich, R. E., 1975. Migration behavior of the larval rodent bot fly *Cuterebra fontinella* in normal and resistant mice. *Folia ent. mexicana* 33: 67-69.
- Gingrich, R. E., 1979. Effects of some factors on the susceptibility of *Peromyscus leucopus* to infestation by larvae of *Cuterebra fontinella* (Diptera: Cuterebridae). *J. Parasitol.* 65 (2): 288-292.
- Gingrich, R. E., 1981. Migratory kinetics of *Cuterebra fontinella* (Diptera: Cuterebridae) in the white-footed mouse, *Peromyscus leucopus*. *J. Parasitol.* 67: 398-402.
- Gingrich, R. E. & C. C. Barrett, 1976. Natural and acquired resistance in rodent hosts to myiasis by *Cuterebra fontinella* (Diptera: Cuterebridae). *J. med. Ent.* 13: 61-65.
- Glasgow, B. J. & J. M. Maggiano, 1995. *Cuterebra* ophthalmomyiasis. *American J. Ophthalmol.* 119: 512-514.
- Glass, E. N., A. M. Cornetta, A. de Lahunta, S. A. Center &

- M. Kent, 1998. Clinical and clinicopathologic features in 11 cats with *Cuterebra* larvae myiasis of the central nervous system. *J. Vet. Intern. Méd.* 12: 365-368.
- Gmelin, J. F., 1788. *Caroli à Linné Systema Naturae, per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus et differentiis. Ed. XIII* 1 (5): 2225-2120. Laur. Salvii, Holmiae [=Stockholm].
- Goddard, J., G. T. Baker & B. R. Norment, 1986. Ectoparasites of the eastern chipmunk (*Tamias striatus*) from Tishomingo County, Mississippi. *J. Wildlife Dis.* 22 ((1): 127-129.
- Goudot, J., 1845. Observations sur un diptère exotique dont la larve nuit aux boeufs (*Cuterebra noxialis*). *Ann. Sci. nat.* 3: 221-230.
- Greene, C. T., 1925. The tentative arrangement of the muscoid flies based on puparia. *Proc. ent. Soc. Washington* 27: 137-163.
- Greene, C. T., 1935. *Cuterebra fontinella*. *Proc. ent. Soc. Washington* 37: 169.
- Guérin-Ménéville, F. E., 1835. Planches des animaux invertébrés. Insectes, pls., in his *Iconographie du règne animal de G. Cuvier, ou représentation d'après nature de l'une des espèces le plus remarquables et souvent non encore figureées de chaque genre d'animaux pouvant servir d'atlas à tous les traités de zoologie*. Vol. 2. Paris.
- Guérin-Ménéville, F. E., 1844. Douzième ordre. Les diptères, pp. 531-559, in his *Iconographie du règne animal de G. Cuvier, ou représentation d'après nature de l'une des espèces le plus remarquables et souvent non encore figureées de chaque genre d'animaux pouvant servir d'atlas à tous les traités de zoologie*. Vol. 3 (Texte explicatif, Insectes), 570 pp. Paris.
- Guimarães, J. H., 1966. Nota sobre os hábitos dos machos de *Dermatobia hominis* (Linnaeus Jr.) (Diptera, Cuterebridae). *Papéis avulsos de Zoologia*, São Paulo 18: 277-279.
- Guimarães, J. H., 1967. Family Cuterebridae, in Papavero, N., ed., *A catalogue of the Diptera of the Americas south of the United States* 105: 1-28. Museu de Zoologia, Universidade de São Paulo, São Paulo.
- Guimarães, J. H., 1971. Notes on the hosts of neotropical Cuterebrini (Diptera, Cuterebridae) with new records from Brazil. *Papéis avulsos de Zoologia*, São Paulo 25 (100): 89-94, figs.
- Guimarães, J. H., 1984. New genera and species of Neotropical Cuterebridae (Diptera, Cyclorrhapha). *Revta bras. Ent.* 28: 23-27.
- Guimarães, J. H., 1990. Revision of the subfamily Rogenhoferinae stat. nov. (Diptera, Cuterebridae). *Mems Inst. Oswaldo Cruz*, Rio de Janeiro 84 (Suppl. IV) ("1989"): 255-261.
- Guimarães, J. H. & C. E. A. Coimbra, 1982. Míase humana por *Alouattamyia baeri* (Shannon & Greene) (Diptera, Cuterebridae). Comunicação de dois casos na Região Norte do Brasil. *Revta bras. Zool.* 11: 35-39.
- Guimarães, J. H. & N. Papavero, 1966. A tentative annotated bibliography of *Dermatobia hominis* (Linnaeus Jr., 1781) (Diptera, Cuterebridae). *Arqos Zool.*, São Paulo 14 (4): 223-294, 2 pls.
- Guimarães, J. H. & N. Papavero, 1999. *Myiasis in man and animals in the Neotropical Region. Bibliographic database*, 308 pp., 120 figs. Fundação de Amparo à Pesquisa do Estado de São Paulo & Editora Plêiade, São Paulo.
- Guimarães, L. R. & M. Carrera, 1941. Contribuição ao conhecimento dos cuterebrídeos do Brasil. *Arqos Zool.*, São Paulo 3 (1): 1-23, 3 pls., 9 figs.
- Harris, T. W., 1833. VIII. Insects, pp. 566-595, in Hitchcock, *Report on the geology, mineralogy, botany, and zoology of Massachusetts*. 1st ed.
- Harris, T. W., 1835. VIII. Insects, pp. 553-602, in Hitchcock, *Report on the geology, mineralogy, botany, and zoology of Massachusetts*. 2nd ed.
- Harris, T. W., 1841. *A report on the insects of Massachusetts, injurious to vegetation*, 459 pp. Cambridge, Massachusetts.
- Haugen, A. O., 1942. Life history studies on the cottontail rabbit in southwestern Michigan. *Amer. Midland Naturalist* 28: 204-244.
- Henriksen, K. L., 1942. A note upon two South American species of warble-flies (*Cuterebra apicalis* Guérin and *schmalzi* Lutz) parasites of rodents. *Ent. Meddr* 22: 284-289, 5 figs.
- Hensley, M. S., 1976. Prevalence of cuterebrid parasitism among woodmice in Virginia. *J. Wildlife Dis.* 12: 172-179.
- Honess, R. F. & K. B. Winter, 1956. Diseases of wildlife in Wyoming. *Wyoming Game Fish Comm. Bull.* 9: 1-279.
- Howship, P., 1833. An account of two cases of inflammatory tumour produced by a deposit of the larva of a large fly (*Oestrus homanus*) beneath the cutis in the human subject; accompanied with drawings of the larva. *Proc. r. Soc. London* 12: 181.
- Hunter, D. M. & J. M. Webster, 1973. Determination of the migratory route of botfly larvaer, *Cuterebra grisea* (Diptera, Cuterebridae) in deer mice. *Intern. J. Parasitol.* 3: 311-316.
- Jacobson, H. A., B. S. McGinnes & E. P. Catts, 1978. Bot fly myiasis of the cottontail rabbit, *Sylvilagus floridanus mallurus* in Virginia, with some biology of the parasite, *Cuterebra buccata*. *J. Wildlife Dis.* 14: 56-66.
- Jellison, W. L., 1949. *Cuterebra thomomuris* sp. nov., a warble from the pocket gopher, *Thomomys talpoides* (Rodentia, Geomyidae). *J. Parasitol.* 35: 482-486.
- Jennison, C. A., L. R. Rodas & G. W. Barrett, 2006. *Cuterebra fontinella* parasitism on *Peromyscus leucopus* and *Ochrotomys nuttalli*. *Southeastern Naturalist* 5 (1): 157-167.
- Johnson, C. W., 1895. Diptera of Florida. *Proc. Acad. nat. Sci. Philadelphia* 1895: 303-340.
- Johnson, C. W., 1903. Diptera of Beulah, New Mexico. *Trans. Amer. ent. Soc.* 29: 101-106.
- Johnson, C. W., 1913. Insects of Florida. I. Diptera. *Bull. Amer. Mus. nat Hist.* 32: 37-90.

- Johnson, C. W., 1925a. Fauna of New England. 15. List of the Diptera or two-winged flies. *Occas. Papers Boston Soc. nat. Hist.* 7: 1-376.
- Johnson, C. W., 1925b. Diptera of the Harris collection. *Proc. Boston Soc. nat. Hist.* 38: 5-99.
- Johnson, C. W., 1930. A bot fly from the white-footed mouse. *Psyche* 37(3): 283-284.
- Joly, N., 1846. Recherches zoologiques, anatomiques, physiologiques et médicales sur les oestrides en général. *Ann. Soc. Agric. Lyon* 9: 157-305, 8 pls. [Also sep. publ., Lyon, 149 pp., 8 pls.].
- Jones, P. R., 1906. A new *Cuterebra* from Nebraska. *Ent. News* 17: 391-392.
- Kemp, D. J. & J. Alcock, 2003. Lifetime resource utilization, flight physiology, and the evolution of contest competition in territorial insects. *American Naturalist* 162(3): 290-301.
- Keth, A. C., 1999. Three incidents of human myiasis by rodent *Cuterebra* (Diptera: Cuterebridae) larvae in a localized region of western Pennsylvania. *J. med. Ent.* 36(3): 831-832.
- Knab, F., 1914. A new *Cuterebra* from Panama. *Insec. Insc. Menstr.* 2: 187-188.
- Knippling, E. F. & A. L. Brody, 1940. Some taxonomic characters of cuterebrine (Diptera) larvae, with larval descriptions of two species from Georgia. *J. Parasitol.* 26: 33-43.
- Knippling, E. F. & W. G. Bruce, 1937. Three unusual records for cuterebrine larvae (Diptera: Oestridae). *Ent. News* 48(6): 156-158.
- Latrelle, P. A., 1803. *In Société de Naturalistes et d'Agriculteurs, Nouveau dictionnaire d'histoire naturelle appliquée aux arts, principalement à l'agriculture et à l'économie rurale et domestique. Nouvelle édition [2ème]* 16. Déterville, Paris.
- Latrelle, P. A., 1818. Oestre, pp. 264-274, in *Société de Naturalistes et d'Agriculteurs, Nouveau dictionnaire d'histoire naturelle appliquée aux arts, principalement à l'agriculture et à l'économie rurale et domestique. Nouvelle édition [2ème]* 23 (NIL-ORC): 612 pp., 6 pls., Déterville, Paris.
- Led, J. E., A. Colacelli, J. J. Boero & E. G. Colombo, 1978. Parasitismo por *Cuterebra apicalis* Guérin, 1829 (Insecta Diptera – Cuterebridae) en rata (*Rattus norvegicus*). *Analecta vet.* 6-8 (1-3): 35-37.
- Led, J. E. & E. Brandelli, 1978. Miasis por *Cuterebra sp.*, Clark, 1815 [sic] (Diptera, Cuterebridae) en una chinchilla (*Chinchilla lanigera*). *Analecta vet.* 6-8 (1-3): 29-33, fig.
- Leite, A. C. R., A. Falquete & M. P. Guimarães, 1985. *Cuterebra schmalzi* Lutz, 1917 (Diptera: Cuterebridae) parasitando *Nectomys squamipes* (Brandt, 1827) (Rodentia: Cricetidae). *Resumos 37º Reunião anual Soc. bras. Progr. Ciência* (Belo Horizonte): 835.
- Leite, A. C. R., M. P. Guimarães & W. S. Lima, 1986. Microscopia eletrônica de varredura do ovo de *Metacuterebra apicalis* (Diptera, Cuterebridae). *Res. XV Congr. bras. Zool.* (Curitiba): 89.
- Leite, A. C. R. & P. Williams, 1988a. The life cycle of *Metacuterebra apicalis* (Diptera, Cuterebridae). *Mems Inst. Oswaldo Cruz*, Rio de Janeiro 83 (4): 485-491.
- Leite, A. C. R. & P. Williams, 1988b. Redescrição de adultos de *Metacuterebra apicalis* (Diptera: Cuterebridae). *Mems Inst. Oswaldo Cruz*, Rio de Janeiro 83: 493-507.
- Leite, A. C. R. & P. Williams, 1989. Morphological observations on the eggs and first instar larvae of *Metacuterebra apicalis* (Diptera: Cuterebridae). *Mems Inst. Oswaldo Cruz*, Rio de Janeiro 84 (1): 123-130.
- Lindquist, A. W., 1937. Myiasis in wild animals in southwestern Texas. *J. econ. Ent.* 30: 735-740.
- Linnaeus, C., Jr., 1781. (Mittheilungen über *Oestrus hominis*). *Pallas' Neue nord. Beytr. Phys. Geogr. Erd. u. Völk.* 1: 157-158.
- Lugger, C., 1896. Insects injurious in 1896. *Bull. Univ. Minnesota agric. Expt Sta., Ent. Div.* 48: 31-270.
- Lugger, C., 1897. Parasites of man and domesticated animals, pp. 44-231, in *2nd annu. Rept Entomologist State Expt Sta. Univ. Minnesota for the year 1896*.
- Lutz, A., 1917. Contribuições ao conhecimento dos oestrideos brasileiros. *Mems Inst. Oswaldo Cruz*, Rio de Janeiro 9: 94-113, pls. 27-29 [Portuguese text in modern orthograph in Bechimol & Sá, 2006: 669-692, pls. 27-29].
- Lutz, A. 1918. A contribution to the knowledge of Brazilian Oestridae. *Mems. Inst. Oswaldo Cruz*, Rio de Janeiro 10: 118-137, pls. 27-29. [Fac-simile reproduction in Bechimol & Sá, 2006: 647-688, pls. 27-29].
- Macquart, J., 1835. *Histoire naturelle des insects. Diptères. Tome deuxième*, 703 pp., 12 pls. Paris.
- Macquart, J., 1843. Diptères nouveaux ou peu connus. Tome deuxième. 3e. partie. *Mém. Soc. Sci. Agric. Arts Lille* 1843: 162-460, 36 pls. [Also sep. publ., pp. 5-304, Paris, 1843].
- Magnarelli, L. A. & T. G. Andreadis, 1981. Human cases of furuncular, traumatic, and nasal myiasis in Connecticut. *American J. trop. Med. Hyg.* 30 (4): 894-896.
- Manrique-Saide, P., S. Hernandez-Betancourt & M. T. Quintero, 2000. First record of *Cuterebra sp.* (Diptera: Cuterebridae) infection in *Otorylomys phyllotis* (Rodentia: Muridae). *Florida Entomologist* 83 (4): 487-488.
- Meigen, J. W., 1824. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten* 4: 428 pp., pls. 33-41. Hamm.
- Mello, D. A., 1978. Nota sobre a biologia de *Cuterebra apicalis* (Guérin 1829/38 [sic]) (Diptera – Cuterebridae). *Revta bras. Pesq. méd. biol.* 11 (6): 329-331.
- Miller, D. H. & L. L. Getz, 1969. Botfly infections in a population of *Peromyscus leucopus*. *J. Mammal.* 50: 277-283.
- Milton, K., 1996. Effects of bot-fly (*Alouattamyia baeri*) parasitism on a free-ranging howler monkey (*Alouatta palliata*) population in Panama. *J. Zool.* 239 (1): 39-63.
- Nelder, M. P. & W. K. Reeves, 2005. Ectoparasites of road-killed vertebrates in northwestern South Carolina, USA. *Vet. Parasitol.* 129: 313-322.

- Noël, S., N. Tessier, B. Angers, D. M. Wood & F. J. Lapointe, 2004. Molecular identification of two species of myiasis-causing *Cuterebra* by multiplex PCR and RFLP. *Med. vet. Ent.* 18 (2): 161-166.
- Olivier, G. A., 1811-[1812]. *Encyclopédie méthodique. Insectes* 8: 1-360 (1811), 361-722 (1812). Paris.
- Painter, R. H., 1930. Notes on Kansas bot-flies (Oestridae: Diptera). *J. Kansas ent. Soc.* 3: 32-35.
- Papavero, N. & S. Ibáñez-Bernal, 2001. Contributions to a history of Mexican Dipterology. – Part I. Entomologists and their work before the *Biología Centrali-Americana. Acta zool. mexicana* (n. s.) 84: 65-173, 10 figs.
- Parker, O. S. & A. H. Chaney, 1979. *Liomys irroratus* (Rodentia: Heteromyidae), a new host for *Cuterebra fontinella* (Diptera: Cuterebridae). *J. med. Ent.* 15: 573.
- Penner, L. R., 1958. Concerning a rabbit cuterebrid, the larvae of which may penetrate the human skin (Diptera, Cuterebridae). *J. Kansas ent. Soc.* 31: 67-71.
- Penner, L. R. & F. P. Pocius, 1956. Nostril entrance as the mode of infection by the first stage larvae of rodent *Cuterebra*. *J. Parasitol.* 42 (4, Sect. 2, Suppl.): 42.
- Pinto, C. M. & G. L. Claps, 2005. First record of *Cuterebra almeidai* (Guimarães and Carrera) from Argentina, new host records for *Cuterebra apicalis* Guérin-Méneville, and list of *Cuterebra* (Diptera: Oestridae) in the collection of the Instituto Fundación Miguel Lillo, Tucumán, Argentina. *Proc. ent. Soc. Washington* 107 (3): 572-575.
- Pfaffenberger, G. S. & V. V. Valencia, 1988. Ectoparasites of sympatric cottontails (*Sylvilagus audubonii* Nelson) and jack rabbits (*Lepus californicus* Mearns) from the High Plains of eastern New Mexico. *J. Parasitol.* 74 (5): 842-846.
- Pruett, J. H. & C. C. Barrett, 1983. Development by the laboratory rodent host of humoral antibody activity to *Cuterebra fontinella* (Diptera: Cuterebridae) larval antigens. *J. med. Ent.* 20: 113-119.
- Pujol-Luz, J. R., A. F. Mendonça & R. Henriques, 2004. Registro de *Gracilinanus agilis* (Marsupialia, Didelphidae) parasitado por *Metacuterebra apicalis* (Diptera, Cuterebridae) no cerrado de Brasília, Distrito Federal, Brasil. *Entomología y Vectores*, Rio de Janeiro 11 (4): 669-672.
- Rice, P. L. & G. E. W. Douglas, 1972. Myiasis in man by *Cuterebra* (Diptera: Cuterebridae). *Ann. ent. Soc. America* 65: 514-515.
- Roberts, R. A., 1933. Additional notes on myiasis in rabbits (Dipt.: Calliphoridae, Sarcophagidae). *Ent. News* 44: 157-159.
- Rodhain, J., 1927. Contribution à la faune des oestridés du Congo Belge. *Ann. Parasitol. hum. comp.* 5: 193-213.
- Rodrigues, M. R., C. B. Weiss & D. W. Muncy, 1974. Ophthalmomyiasis of the eyelid caused by *Cuterebra* larva. *Amer. J. Ophthalmol.* 78: 1024-1026.
- Sabrosky, C. W., 1972. Discovery of the bot fly collection of Bracy Clark, with designation of lectotypes in *Cuterebra*. *J. Ent. (Ser. B, Taxon. Syst.)* 41: 89-96.
- Sabrosky, C. W., 1986. *North American species of Cuterebra. The rabbit and rodent bot flies (Diptera: Cuterebridae)*, 240 pp. Thomas Say Foundation, Ent. Soc. America, College Park, Maryland [Vol. 11].
- Scaglia, O. A. & C. A. Velázquez, 1982. Incidencia de la parasitosis producida por larvas de *Rogenhofera* sp. (Diptera: Cuterebridae) en una comunidad de roedores en el sudeste de la provincia de Buenos Aires. *Res. X. Reun. argent. Ecol.* (Mar del Plata).
- Schiff, T. A., 1993. Furuncular cutaneous myiasis caused by *Cuterebra* larva. *J. American Acad. Dermatol.* 28: 261-263.
- Scholten, T. H. & V. H. Chrom, 1979. Myiasis due to *Cuterebra* in humans. *J. Canadian. med. Ass.* 120: 1392-1393.
- Scott, T. G. & E. Snead, 1942. Warbles in *Peromyscus leucopus novaeboracensis*. *J. Mammal.* 23 (1): 94-95.
- Shannon, R. C., 1925. (*Cuterebra* from howling monkey of tropical America; abstr. of report before Ent. Soc. of Washington). *J. Washington Acad. Sci.* 15 (15): 379.
- Shannon, R. C. & E. Del Ponte, 1926. Sinopsis parcial de los muscoideos argentinos. *Revta Inst. Bacteriol.*, Buenos Aires 5: 1-44, 4 pls.
- Shannon, R. C. & C. T. Greene, 1956. A bot-fly parasite in monkeys. *Zoopathologica* 1 (7): 287-290, figs. 152-153, pls. A-B.
- Shiffer, C. N., 1983. Aggregation behavior of adult *Cuterebra fontinella* (Diptera: Cuterebridae) in Pennsylvania, USA. *J. med. Ent.* 20: 365-370.
- Shoemaker, J. P. & S. J. Joy, 1968. Some ectoparasites from West Virginia mammals. *Proc. West Virginia Acad. Sci.* (1967) 39: 78-80.
- Sillman, E. I., 1956. Studies on the biology of a cuterebrid (Cuterebridae: Diptera) infesting *Peromyscus leucopus novaeboracensis*, Fischer, the white-footed mouse, in southern Ontario. *Annu. Rept ent. Soc. Ontario* (1955) 86: 89-97.
- Sillman, E. I., 1957a. Further laboratory and field observations on the ecology of some Ontario Cuterebridae (Diptera), in particular, *Cuterebra angustifrons* Dalmat 1942. *Annu. Rept ent. Soc. Ontario* (1956) 87: 28-40.
- Sillman, E. I., 1957b. Some aspects of the biology of *Cuterebra angustifrons* Dalmat, 1942 (Cuterebridae, Diptera). *J. Parasitol.* 43: 33-34.
- Sillman, E. J. & M. V. Smith, 1959. Experimental infestation in *Peromyscus leucopus* with larvae of *Cuterebra angustifrons*. *Science*, New York 130 (3368): 165-166.
- Slansky, F., 2006. *Cuterebra* bot flies (Diptera: Oestridae) and their indigenous hosts and potential hosts in Florida. *Florida Entomologist* 89 (2): 152-160.
- Slansky, F. & J. Huckabee, 2006. First record of rodent-infesting *Cuterebra* bot flies parasitizing raccoons (*Procyon lotor*) in North America. *J. Parasitol.* 92 (6): 1369-1373.
- Smith, J. B., 1900. *Insects of New Jersey*, 755 pp. New Jersey State Board of Agriculture, Trenton.
- Smith, J. B., 1910. The insects of New Jersey. *Annu. Rept. New Jersey State Mus.* 1909: 1-888. MacCrellish and Quigley, Trenton.

- Smith, R. C., 1943. Common insects of Kansas. *Rept. Kansas State Bd. Agric.*, 40 pp.
- Stuart, M., V. Pendergast, S. Rumfelt, S. Pierberg, L. Greenspan, K. Glander & M. Clarke, 1998. Parasites of wild howlers (*Alouatta spp.*). *Intern. J. Primatology* 19 (3): 493-512.
- Swenk, M. H., 1905. The North American species of *Cuterebra*. *J. New York ent. Soc.* 13: 181-185.
- Timm, R. M. & E. F. Cook, 1979. The effect of bot fly larvae on reproduction in white-footed mice, *Peromyscus leucopus*. *American Midland Naturalist* 101: 211-217.
- Timm, R. M. & R. E. Lee, Jr., 1981. Do bot flies, *Cuterebra* (Diptera: Cuterebridae), emasculate their hosts? *J. med. Ent.* 18 (4): 333-336.
- Townsend, C. H. T., 1892. Descriptions of oestrid larvae taken from the jack rabbit and cotton-tail. *Psyche* 6: 298-300.
- Townsend, C. H. T., 1893. Further notes on the cottontail bot, with the breeding and identification of the fly. *Insect Life* 5: 317-320.
- Townsend, C. H. T., 1895a. Contributions to the dipterology of North America. II. Tabanidae, Conopidae, Tachinidae, etc. *Trans. American ent. Soc.* 22: 55-80.
- Townsend, C. H. T., 1895b. Notes on the Diptera of Baja California, including some species from adjacent regions. *Proc. California Acad. Sci.* (2) 4: 593-620.
- Townsend, C. H. T., 1897. Description of the bot-fly of the cotton-tail rabbit in New Mexico, *Cuterebra lepusculini* sp. *Psyche* 8: 8-9.
- Townsend, C. H. T., 1911a. Announcement of further results secured in the study of muscoid flies. *Ann. ent. Soc. America* 4: 127-152.
- Townsend, C. H. T., 1911b. Correction to my paper in June (1911) issue of the Annals E. S. A. *Ann. ent. Soc. America* 4: 328-329.
- Townsend, C. H. T., 1913. Inquiry into the relationships and taxonomy of the muscoid flies. *Canadian Entomologist* 45: 37-57.
- Townsend, C. H. T., 1917. A synoptic revision of the Cuterebridae, with synonymic notes and description of one new species. *Insec. Insc. Menstr.* 5: 23-28.
- Townsend, C. H. T., 1919. New genera and species of muscoid flies. *Proc. U. S. natl Mus.* 56: 541-592.
- Townsend, C. H. T., 1927. Synopse dos generos muscoideos da região humida tropical da América, com generos e espécies novas. *Revta Mus. paulista* 15: 205-385, 7 figs.
- Townsend, C. H. T., 1931. New genera and species of American oestromuscid flies. *Revta Ent.*, Rio de Janeiro 1: 313-354.
- Townsend, C. H. T., 1935. *Manual of myiology* 2: 289 pp., 9 pls. Itaquaquecetuba, São Paulo.
- Townsend, C. H. T., 1938. *Manual of myiology* 6: 142 pp. Itaquaquecetuba, São Paulo.
- Townsend, C. H. T., 1942. *Manual of myiology* 12: 342 pp., 84 pls. Itaquaquecetuba, São Paulo. [In Addenda & corrigenda, p. 333].
- Vieira, E. M., 1993. Occurrence and prevalence of the bot flies *Metacuterebra apicalis* (Diptera: Cuterebridae), in rodents of Cerrado from Central Brazil. *J. Parasitol.* 79(5): 792-795.
- Vignau, M. L. & G. A. Zuleta, 1991. Descripción de los estadios preimaginales de *Rogenhofera bonaerensis* (Del Ponte, 1939) (Diptera, Cuterebridae). *Stud. neot. Fauna Environm.* 26 (3): 129-133.
- Walker, F., 1849. *List of the specimens of dipterous insects in the collection of the British Museum* 3: 485-687. London.
- Washburn, F. L., 1905. The Diptera of Minnesota. *Bull. Univ. Minnesota agric. expt Sta.* 93: 19-168.
- Weisbroth, S. H., R. Wang & S. Scher, 1973. *Cuterebra buccata*: Immune response in myiasis of domestic rabbits. *Exptl Parasitol.* 34 (1): 22-31.
- Wiedemann, C. R. W., 1830. *Aussereuropäische zweiflügelige Insekten* 2: xii + 684 pp., 5 pls., Hamm.
- Williston, S. W., 1884. Order VII. Diptera, pp. 403-432, 433, figs. 508-594, in Kingsley, J. S., ed., *The standard natural history* 2: 555 pp., 666 figs., 20 pls. Boston.
- Wilson, W. D., J. A. Hnida & D. W. Duszynski, 1997. Parasites of mammals on the Sevilleta National Wildlife Refuge, Socorro, New Mexico: *Cuterebra austeni* and *C. neomexicana* (Diptera: Oestridae) from *Neotoma* and *Peromyscus* (Rodentia: Muridae), 1991-1994. *J. med. Ent.* 34 ((3)): 359-367.
- Wolf, M. & G. O. Batzli, 2001. Increased prevalence of bot flies (*Cuterebra fontinella*) on white-footed mice (*Peromyscus leucopus*) near forest edges. *Canadian J. Zool.* 79 (1): 106-109.
- Wulp, F. M. van der, 1888. Family Oestridae, pp. 1-2, in Godman, F. D. & O. Salvin, eds. *Biologia-Centrali-America. Zoologia-Insecta-Diptera* 2: 1-489. London.
- Zeledón, R., O. Jiménez Q. & R. R. Brenes M., 1957. *Cuterebra baeri* Shannon & Greene en el mono aullador de Costa Rica. *Revta Biol. trop.*, San José 5: 129-134, 11 figs.
- Zimšen, E., 1964. *The type-material of I. C. Fabricius*, 656 pp. Munksgaard, Copenhagen.
- Zuleta, G. A. & M. L. Vignau, 1990. Bot fly parasitism (*Rogenhofera bonaerensis*) (Diptera, Cuterebridae) in the pampean grassland mouse (*Akodon azarae*), in Argentina. *J. Wildlife Dis.* 26 (1): 11-17.

Index

- albifrons* Swenk, 1905, Cuterebra – 4
albipilosa Sabrosky, 1986, Cuterebra – 2
almeidai (L. R. Guimarães & Carrera), 1941 (Baucuterebra), Metacuterebra – 5
analis (Macquart), 1843 (Cuterebra), Metacuterebra – 5
Alouattamyia Townsend, 1931 – 5
americana Fabricius of Johnson, 1903, Cuterebra, misident. – 3
americana Fabricius of Walker, 1849, Cuterebra, misident. – 4
americana Fabricius of Wulp, 1888, Cuterebra, part, misident. – 5
analis (Macquart), 1843 (*Cuterebra*), Metacuterebra – 5
Andinocuterebra J. H. Guimarães, 1984 – 2
angustifrons Dalmat, 1942, Cuterebra – 4
apicalis (Guérin-Ménéville), 1835 (*Cuterebra*), Metacuterebra – 5
approximata Walker of Wulp, 1888, Cuterebra, misident. – 4
arizonae Sabrosky, 1986, Cuterebra – 2
atrox Clark, 1849, Cuterebra – 2
Atrypoderma Townsend, 1919 – 2
Atryposoma Bau, 1931, error – 2
austeni Sabrosky, 1986, Cuterebra – 3

bacata Washburn, 1905, Cuterebra, error – 3
baccatta Lugger, 1896, 1897, Cuterebra, error – 3
baeri (Greene), 1925 (*Cuterebra*), Metacuterebra – 5
bajensis Sabrosky, 1986, Cuterebra – 3
Baucuterebra L. R. Guimarães & Carrera, 1941 – 5
baui (Fonseca), 1940 (*Oricuterebra*), Metacuterebra – 6
bicincta (Bau), 1931 (Cuterebra; as *versicolora* spp.), Metacuterebra – 6
Boegeria Austen, 1895 – 2
bonaerensis (Del Ponte), 1939 (Guerinioestrus), Rogenhofera – 8
buccata (Fabricius), 1777 (*Oestrus*), Cuterebra – 3
buccata Fabricius of Baker, 1904, Cuterebra – 4
buccata Fabricius of Johnson, 1913, Cuterebra, misident. – 4
buccator Williston, 1884, Cuterebra, error – 3
buccatta Eddy & Emerson, 1940, Cuterebra, error – 3
bureni (Dalmat), 1942 (*Montemyia*), Metacuterebra – 6

cayennensis (Macquart), 1843 (Cuterebra), Metacuterebra – 6
clarki Sabrosky, 1986, Cuterebra – 3
cometes (Shannon & Del Ponte), 1926 (Cuterebra), Metacuterebra – 5
conflaps (Bau), 1929 (Cuterebra), Metacuterebra – 6
cuccatus Harris, 1833, Cuterebra, error – 3
?cuniculi Clark of Harris, 1833 and Smith, 1943, Cuterebra, misident. – 3
cuniculi Clark of Johnson, 1895, Cuterebra, misident. – 4
Curtebra Blanchard, 1840, error – 2
Curteterbra Blanchard, 1845: 484, error – 2
Cuterebra Clark, 1815 – 2
Cuterebrella J. H. Guimarães, 1984 – 2

Cuterebridae, Brauer, 1887 – 1
Cuterebriden (vernacular) Brauer, 1863 – 1
Cuterebrinae Bau, 1906 – 1
Cuterebrine Townsend, 1913 – 1
Cuteribrinae Rodhain, 1927 – 2
Cutirebra Wulp, 1888, error – 2
Cutiterebra Agassiz, 1846, emend. – 2
Cutiterebrinae Austen, 1895 – 1
cyaniventris (Macquart), 1843 (Cuterebra), Dermatobia – 5

daguerrei Del Ponte, 1939 (Guerinioestrus), Rogenhofera – 8
dasypoda Brauer, 1896, Rogenhofera – 8
Dermatobia Brauer, 1861 – 5
Dermatobiidae Bromley, 1937 – 2
Dermatobiiae Townsend, 1913 – 1
Dermatobiinae Townsend, 1917 – 2
detrudatrix (Clark), 1848 (Cuterebra), Metacuterebra – 6
dorsalis (Bau), 1929 (Cuterebra), Metacuterebra – 6

elegans (Bau), 1931 (Cuterebra, as *rufiventris* spp.), Metacuterebra – 7
ephippium Latreille, 1818 (Cuterebra) – 7

fasciata Swenk of Townsend, 1917, Cuterebra, misident. – 4
fassleri J. H. Guimarães, 1984, Andinocuterebra – 2
flaviventris Bau, 1931 (Metacuterebra) – 7
fontanella Clark, 1843, Cuterebra, error – 4
fontinella Clark, 1827, Cuterebra – 3
fontinella Clark of Townsend, 1893, 1895 and Honess & Winter, 1956, Cuterebra, misident. – 4
frontinella Fletcher & Gibson, 1908, Cuterebra, error – 4
funebris (Austen), 1895 (Cuterebra), Metacuterebra – 6

gilvopilosa Bau, 1932, Rogenhofera – 8
grandis (Guérin-Ménéville), 1844 (*Cephenemyia*), Rogenhofera – 8
Guerinioestrus Townsend, 1931 – 8

hermanni Bischoff, 1900, Pseudogametes – 7
histrio Coquillett, 1902, Cuterebra – 4
hominis (Linnaeus Jr.), 1781 (*Oestrus*), Dermatobia – 5
horripilum Clark of Walker, 1849, Cuterebra, misident. – 3
humanus (Howship), 1833 (*Oestrus*), Dermatobia – 5

infulata (Lutz), 1917 (Cuterebra), Metacuterebra – 6

jorgensenii Del Ponte, 1959, Metacuterebra, *nomen nudum* – 8

lepidia (Austen), 1933 (Cuterebra), Metacuterebra – 6
lepusculi Townsend, 1897, Cuterebra – 4
lopesi J. H. Guimarães, 1989, Rogenhofera – 8
lutzi Bau, 1930 (Cuterebra) – 7

- maculosa Knab, 1914, Cuterebra – 4
 megastoma (Brauer), 1863 (Cuterebra), Metacuterebra – 6
 Metacuterebra Bau, 1929 – 5
Montemyia Dalmat, 1942 – 5
- nigricans* (Lutz), 1917 (Cuterebra), Metacuterebra – 6
nigricincta (Austen), 1895 (Cuterebra), Metacuterebra – 6
noxialis (Goudot), 1845 (Cuterebra), Derematobia – 5
- obidosiana* (Bau), 1929 (Cuterebra, as *conflans* ssp.),
 Metacuterebra – 6
Oricuterebra Bau, 1931 – 5
ornata Bau, 1928 (Cuterebra) – 7
Orthocuterebra Bau, 1929 – 2
- Paracuterebra* Bau, 1929 – 2
patagona (Guérin–Ménéville), 1844 (Cuterebra),
 Metacuterebra – 6
peromysci Dalmat, 1942, Cuterebra – 4
pessoai (L. R. Guimarães & Carrera), 1941 (Cuterebra),
 Metacuterebra – 7
postica Sabrosky, 1986, Cuterebra – 4
praegrandis Austen, 1933, Cuterebra – 4
princeps (austen), 1895 (*Bogeria*), Cuterebra – 4
Protocuterebra Bau, 1929 – 2
Pseudobogeria Bau, 1931 – 2
Pseudogametes Bischoff, 1900 – 7
Pseudogametinae Townsend, 1917 – 2
Pseudogametine [sic] Townsend, 1911 – 1
Pseudogametini Townsend, 1911 – 1
Pseudomagetini Dalmat, 1942, error – 2
pulchra (Bau), 1930 (Cuterebra), Metacuterebra – 6
purivora Clark, 1815, Cuterebra – 3
pusilla (Bau), 1931 (Cuterebra), Metacuterebra – 5
pygmaea (Bau), 1931 (Cuterebra), Metacuterebra – 6
- ribeiroi* (Artigas & d'Andretta Jr.), 1951 (*Alouattamyia*),
 Metacuterebra – 6
Rogenhofera Brauer, 1863 – 7
- Rogenhoferia*, error or emend. – 8
Rogenhoferinae J. H. Guimarães, 1990 – 2
rubiginosa (Bau), 1931 (Cuterebra), Metacuterebra – 6
rufiventris (Macquart), 1843 (Cuterebra), Metacuterebra – 7
- sabroskyi* (J. H. Guimarães), 1984 (Cuterebrella), Cuterebra – 4
sarcophagooides (Lutz), 1917 (Cuterebra), Metacuterebra – 6
schmalzi (Lutz), 1917 (Cuterebra), Metacuterebra – 7
schroederi (Enderlein), 1909 (Cuterebra), Metacuterebra – 6
scudderii Townsend, 1917, Cuterebra – 3
semiater (Wiedemann), 1830 (Musca), Pseudogametes – 7
semilutea Bau, 1929 (Cuterebra) – 7
similis Johnson, 1903, Cuterebra – 3
simulans (Austen), 1933 (Cuterebra), Metacuterebra – 7
sp. Adler, Davis & Carvajal, 2003 (Cuterebra) – 7
sp. Austen, 1895, Cuterebra – 4
sp. Curran, 1934, Cuterebra – 3
sp. Led & Brandelli, 1978 (Cuterebra) – 7
sp. Manrique-Saide, Hernandez-Betancourt & Quintego, 2000, Cuterebra – 7
sp. Sillman, 1956, Cuterebra – 4
sp. Townsend, 1892, Cuterebra – 4
sp. Townsend, 1911, Cuterebra – 4
sp. near albipilosa Sabrosky, 1986, Cuterebra – 4
subbuccata Bau, 1929, Cuterebra – 4
- terrisona* Walker, 1849, Cuterebra – 4
townsendi (Fonseca), 1941 (*Oricuterebra*), Metacuterebra – 7
trigonocephala Lutz, 1917, *Rogenhofera*, error – 8
trigonophora Brauer, 1863, *Rogenhofera* – 8
Trypoderma Wiedemann in Meigen, 1824 – 2
- versicolora* (Bau), 1939 (Cuterebra), Metacuterebra – 6
vianai Del Ponte, 1959, *Guerinioestrus*, *nomen nudum* – 8
- worontzowi* L. R. Guimarães & Carrera, 1941 (Cuterebra),
 Metacuterebra – 6
- Xylocopodes* Townsend, 1927 – 7

